NOV 26, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.8zshx6e

Protocol Citation: Srinidhi Holalu, Angela Fang, Benjamin Blackman 2019. *In vitro* testing of guide RNA efficiency for CRISPR-mediated genome editing. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.8zshx6e>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Nov 05, 2019

Last Modified: Nov 26, 2019

PROTOCOL integer ID:
 29458

Keywords: gRNA, gRNA testing, CRISPR, gRNA efficiency, genome editing, *In vitro*

In vitro testing of guide RNA efficiency for CRISPR-mediated genome editing

Srinidhi Holalu¹, Angela Fang¹, Benjamin Blackman¹

¹Department of Plant and Microbial Biology, University of California, Berkeley

Mimulus

Benjamin Blackman
 University of California, Berkeley

ABSTRACT

In CRISPR-mediated genome editing, guide RNAs (gRNAs) serve to target the Cas9 nuclease to an intended target genomic locus. Many gRNAs are not efficient in facilitating Cas9-mediated cleavage by introduction of double-strand breaks. Therefore, after designing possible gRNA sequences via web-based tools, it is critical to assess their cleavage efficiency at the target locus and identify the most efficient gRNA(s) *in vitro* before deploying in binary plasmids for genome editing *in planta*.

ATTACHMENTS

[Invitro_test_gRNA_090419.pdf](#)

MATERIALS

MATERIALS

- HiScribe T7 Quick High Yield RNA Synthesis Kit - 50 rxns **New England Biolabs Catalog #E2050S**
- Cas9 Nuclease, *S. pyogenes* - 70 pmol **New England Biolabs Catalog #M0386S**

Reagents necessary:

- Zymo/Bioneer DNA cleanup kit
- CTAB extraction buffer for genomic DNA extraction
- Nuclease free water
- Equipment and reagents to run DNA products through standard agarose gel electrophoresis
- Phusion DNA polymerase kit or any high-fidelity DNA polymerase

For safety information and warnings, please refer to the SDS (Safety Data Sheet).

Cloning a template fragment from genomic DNA

- 1 Perform DNA extraction using CTAB method or any other methods that you may use routinely for your plant species.
- 2 Design primers flanking your target PAM sites (NGG). PCR to amplify around 500-600 bp of genomic/gene fragment around the target. Large size templates are easy to resolve on gel electrophoresis.
- 3 Clean PCR product using Zymo/Bioneer DNA cleanup kit and Sanger sequence to confirm that your PCR amplifications are intended gene. This will also help later for screening indel polymorphisms at PAM site.

Designing gRNA and synthesis of guide RNA using *in vitro* T..

- 4 Design targets for editing target exon(s) using gRNA design tool (<http://crispr.mit.edu/> or any other portal). The following primers are then used to synthesize the DNA template to prepare gRNA for each target.

Oligo design

a. Primer 1 (specific for each guide):

a) TAATACGACTCCTATAGGG-(20 bp guide sequence)-gtttAagagctaTGCTGgaa

b) The three bolded G's are included in the transcript and required for transcription. If the guide-specific sequence starts with one or more G's, the same number of bolded G's can be deleted from the primer.

b. Primer 2 (always the same):

a) Reverse-gRNA Template:

aaaagcaccgactcggtgccacttttcaagttgataacggactagccttatttAaacttgctaTGCTGttc
CAGCAtagctctTaaac

Designing gRNA and synthesis of guide RNA using *in vitro* T..

5

Set up the following PCR reaction:

- 30.5 μL nuclease free water
- 10 μL 5X Phusion buffer
- 1 μL 10 millimolar (mM) dNTP
- 4 μL primer 1 (10 micromolar (μM))
- 4 μL primer 2 (10 micromolar (μM)) Always Primer 2 (Reverse-gRNA Template)
- 0.5 μL Phusion polymerase

Total 50 μL

6

Program for PCR:

1. 98 $^{\circ}\text{C}$ 00:01:00
2. 98 $^{\circ}\text{C}$ 00:00:30
3. 55 $^{\circ}\text{C}$ 00:00:15

8X-10X cycles of these three steps

4. 72 $^{\circ}\text{C}$ 00:01:00
5. 72 $^{\circ}\text{C}$ 00:05:00
6. 4 $^{\circ}\text{C}$ hold

7

Nano-drop to quantify the PCR product. Run a small aliquot of the PCR product out on a gel to check the bands. The resulting band should be **90 bp** (very small) and so it is necessary to run a **$\geq 3\%$ agarose gel**.

8

Purify the PCR product using Zymo/bioneer DNA cleanup kit.

gRNA synthesis

9

Note

Use HiScribe T7 High Yield RNA Synthesis Kit (NEB E2050S)

10

Set up reaction in PCR tube:

- 7 μL water (nuclease free)
- 5 μL NTP mix [10 millimolar (mM) each
- 2 μL Template DNA (~ 100 ng – 500 ng)
- 1 μL T7 RNA Polymerase Mix

Total 15 μL

11 Incubate at 37 °C for 02:00:00 .

12 Move the reaction into 1.5 mL eppendorf tube.

13 Add 435 μL DNAase/RNase free water.

14 Measure gRNA concentration on nanodrop. Note it down. (May not be accurate, as NTPs may inflate the nanodrop readings).

15 Optional to cleanup using any RNA clean-up kit. Dilute to 10 ng / μL by adding nuclease free water.

Cas9 nuclease assay and gel electrophoresis: Incubation of..

16 Set up reaction (include a negative RNA control):

- 7.5 μL water (nuclease free)
- 1.5 μL 10X Cas9 reaction buffer
- 1.5 μL 300 nanomolar (nM) sgRNA (10 ng / μL)
- 0.5 μL Cas9 Nuclease

Total 11 μL

17 Pre-incubate for 00:10:00 at 37 °C .

18 Add 4 μ L substrate DNA (~ 50 ng – 100 ng total).

19 Incubate 01:00:00 at 37 °C (or longer up to 01:30:00).

20 Denature Cas9 by incubation at 80 °C for 00:10:00 .

21 Set up a control reaction including all the above ingredients without gRNA. In addition, different ratios of gRNA to Cas9 nuclease can also be setup to replicate the efficiency tests.

Cas9 nuclease assay and gel electrophoresis: Gel Electroph..

22 Prepare 1.1 % volume agarose gel.

23 Mix the Cas9 incubated reaction with 2 μ L of loading dye. Load the mixture on gel. If stain was not added during gel preparation, soak gel in staining buffer.

24 Run gel electrophoresis until the ladder resolves and check the cleavage of template DNA under UV illumination.