AUG 12, 2020

🌐 Indirect ELISA for detection of anti-HIV antibodies in cats immunized with a HIV gp120 peptide.

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bjnpkmdn

Protocol Citation: Angel A Justiz-Vaillant 2020. Indirect ELISA for detection of anti-HIV antibodies in cats immunized with a HIV gp120 peptide..

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bjnpkmdn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Aug 12, 2020

Last Modified: Aug 12, 2020

PROTOCOL integer ID:
40367

- 1 The ELISA describes detection of antibodies against HIV proteins. It is an ELISA with a modification. The plate is coated with HIV peptides, fragment 254-274 of gp120.

- 2 The modification is that triplicates of 1/16 dilutions of cat sera in PBS non-fat milk are added to the micro-wells.
- 3 After incubation for 90 min at RT the microplate is washed 4X using PBS-Tween 20.
- 4 Then, 25 µl of recombinant protein LA-HRP diluted 1:1000 (Sigma-Aldrich) is added to the assay.
- 5 The microplate is incubated for 1h at RT and then, it is washed four times.
- 6 Tetramethylbenzidine (TMB) solution (50 µl) is added to each well. .
- 7 After a further incubation of 15 min in the dark, the reaction is stopped.
- 8 The microplate is read at 450 nm in a microplate reader at 450 nm. Geometric mean antibody titer is calculated using Perkins method.