AUG 12, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bjnqkmdw

Protocol Citation: Angel A Justiz-Vaillant 2020. Inhibition immunoassay of the HIVgp120_anti-HIVgp120 (Ab-1) reaction by anti-anti-idiotypic-HIVgp120 antibody (Ab-3). . **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjnqkmdw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 12, 2020

Last Modified: Aug 12, 2020

PROTOCOL integer ID:
 40368

Inhibition immunoassay of the HIVgp120_anti-HIVgp120 (Ab-1) reaction by anti-anti-idiotypic-HIVgp120 antibody (Ab-3).

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

MATERIALS

MATERIALS

☒ Carbonate Coating Buffer **BioLegend Catalog #421701**

☒ Anti-Chicken IgY, HRP Conjugate, 300ul **Promega Catalog #G1351**

☒ PBS Blocking buffer with non fat milk **Bio Basic Inc. Catalog #SB0625.SIZE.1pk**

☒ 96-Well Microtiter™ Microplates, Polystyrene, 400μL, Nonsterile, U-bottom **ThermoFisher Catalog #2205**

☒ TMB Substrate Set **BioLegend Catalog #421101**

☒ Fragment 254-274 of Gp120 of HIV (peptide) **Contributed by users**

- 1 Ninety six well polystyrene microplate (U-shaped bottom, Sigma-Aldrich Co, St Louis Missouri) is coated with 50 μl/well of 1 ng/μl solution of gp-120 peptide, fragment 254-274 (Sigma) in

carbonate-bicarbonate buffer pH 9.6 (Sigma) for 4 hours at 37°C.

- 2 The microplate is then washed 4 times with PBS Tween-20.
- 3 The microplate is blocked with 3% non-fat milk in PBS, 25 µl/well, 1h, RT.
- 4 Triplicates of serial doubling dilutions of cat serum diluted in PBS-3% non-fat milk, pH 7.4 are added to the microplate which is then incubated for 90 min at 37°C.
- 5 The microplate is washed 4 times with 150 µl/well of PBS-Tween 20, and 25 µl of a water soluble fraction (WSF) containing anti-HIV-gp120 antibodies (titer =1:1024) is added to each well.
- 6 After adding WSF and it follows an incubation period (90 min, 37°C).
- 7 The microplate is washed 4 times and 25 µl of a 1:30,000 dilution of rabbit anti-chicken IgY-HRP (Sigma-Aldrich Co) is added to each well.
- 8 The microplate is then incubated at 37°C for 1 h.
- 9 The microplate is washed with PBS-Tween 20, 4 times.

- 10 Add 25 µl of TMB to the each well of the microplate and incubate the plate in the dark for 15 min.
- 11 The reaction was stopped with 3M H₂SO₄.
- 12 The microplate is read at 450 nm in a microplate reader.