VERSION 1

FEB 01, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.9dph25n

Protocol Citation: Jamie Allen, Elizabeth Neumann, Maya Brewer, Haichun Yang, Jeff Spraggins, Danielle Gutierrez 2020. Initial Rapid Pathology Assessment of Kidney Tissue. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.9dph25n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Nov 15, 2019

Last Modified: Apr 08, 2023

PROTOCOL integer ID:
29839

Keywords: HuBMAP, Kidney, Quality Assessment, PAS

Initial Rapid Pathology Assessment of Kidney Tissue V.1

Jamie Allen¹, Elizabeth Neumann¹, Maya Brewer¹, Haichun Yang¹, Jeff Spraggins¹, Danielle Gutierrez¹

¹Vanderbilt University

VU Biomolecular Multimodal Imaging Center / Spraggins Research Group

Human BioMolecular Atlas Program (HuBMAP) Method Development

Jamie Allen
Vanderbilt University

ABSTRACT

Scope:

Prepare formalin fixed tissue with freezing fresh tissue protocol (dx.doi.org/10.17504/protocols.io.6wghfbw).

This protocol provides the steps to prepare formalin fixed human kidney tissue and perform histology assessment for normalcy.

Expected Outcome:

Pathological assessment of kidney tissue for use in LC-MS/MS and imaging.

- 1 Once fixed (dx.doi.org/10.17504/protocols.io.6wghfbw) take tissue to pathology Core for paraffin emedding.
- 2 Section samples at 5 µm.
- 3 PAS stain tissue sections (dx.doi.org/10.17504/protocols.io.4qngvve).
- 4 Scan slides with brightfield scanner (Leica) and save as .tiff or .jpg
- 5 Place saved images on QuPath for analysis.

6 Assess and record the following information for each tissue:
(%): cortex and medulla
(Yes/No): Pyramid Presence, Autolysis, Non-Renal disease (*i.e.* cancer)
(0=None – 3=Severe): Glomerular disease, Tubulointerstitial disease

7 Based on FFPE assessment, kidneys will be used accordingly:

LC-MS/MS normalcy:

low or no glomerular disease, and 75% or more cortex for normal
100% tumor for diseased

3D Imaging:

50:50 cortex:medulla

low or no glomerular disease

no renal disease