AUG 22, 2020

Isolation of Salmonella pathogens from oysters

Sade Aisha Folashade

John¹,

Patrick E. Akpaka¹,

Chandrashekhar Unakal¹, Arvind Kurhade¹, Angel Justiz-Vaillant²

¹University of the West Indies St. Augustine;

²University of the West Indies. St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

OPEN ACCESS

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

DOI:

dx.doi.org/10.17504/protocols.io.bj6qkrdw

Protocol Citation: Sade Aisha Folashade John, Patrick E. Akpaka, Chandrashekhar Unakal, Arvind Kurhade, Angel Justiz-Vaillant 2020. Isolation of Salmonella pathogens from oysters.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bj6qkrdw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Aug 22, 2020

Last Modified: Aug 22, 2020

PROTOCOL integer ID:
40880

1 The filtered homogenate was streaked using standardized loops on Salmonella Shigella agar

(selective and differential media for Salmonella and Shigella); the loops were flamed periodically to ensure sterility. This was done in duplicate. The plates were then incubated at 34°C. After an overnight incubation at 34°C, the plate with the best significant and adequate colonies was used to test for the presence of Salmonella. Those colonies that were morphologically characteristic for Salmonella are gram stained:

- 2 Each slide was labelled according to location and given a number. A circle using a wax pencil was drawn on the underside of each slide.
- 3 Drop of sterile saline solution was placed into the circle and using a sterile loop, a single colony from the plate was taken and mixed into it.
- 4 The smear was allowed to air dry and then heat fixed by holding the slide on one side with a sterile forcep and passing the entire slide through the flame of a Bunsen burner two or three times with the smear side up. Making sure that overheating does occur.
- 5 The heat fixed slide was placed on a straining tray.
- 6 The slide was gently flood with crystal violet and left to stand for 1 minute.
- 7 Using a wash bottle of distilled water, the slide was tipped slightly and gently rinsed.
- 8 Then using Gram's iodine, the slide was gently flood and let stand for 1 minute.
- 9 The slide was tipped slightly and gently rinsed with distilled water using a wash bottle.

- 10 95% ethyl alcohol was used to decolorize the smear by for 5-10 seconds.
- 11 The slide was then immediately rinsed with water.
- 12 Using Safranin, the slide was gently flood to counterstain and let stand for 45 seconds.
- 13 The slide was tipped slightly and gently rinsed with distilled was using a wash bottle.
- 14 The slide was then blot dry ice with bibulous paper.
- 15 Using a light microscope under oil immersion, the smear was observed to the determine whether the colonies were gram negative or gram positive.
- 16 This was done per presumptive Salmonella colony observed on each plate.
- 17 A single colony of presumptive Salmonella was then subcultured on SS agar and incubated at 34°C to purify the culture. A single colony was later gram stained prior to biochemical testing in order to ensure the culture's purity. The pure single colonies were further analysed via

biochemical tests (Indole, Citrate, Urease, Motility and Triple Sugar Iron Agar) together with a control *Salmonella* colony taken from the laboratory in order to make a comparison. A single colony of the organism was then saved in a cryovial containing Nutrient Agar at -70 °C. This was done per *Salmonella* spp. isolate found.