OCT 14, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.768hrhw

Protocol Citation: Igem Dusseldorf 2019. Isolation of total RNA from Synechocystis. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.768hrhw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 14, 2019

Last Modified: Oct 14, 2019

PROTOCOL integer ID:
 28608

Isolation of total RNA from Synechocystis

Igem Dusseldorf¹

¹Heinrich-Heine Universität Düsseldorf

Igem Dusseldorf

ABSTRACT

Isolation of total RNA from Synechocystis sp. PCC 6803 Can be used for further work, like qPCR Always work fast, with ice and RNase-free (wear gloves!)

- PGTX solution (100 mL)
- 39.6 g phenol, 6.9 mL glycerol, 0.1 g hydroxyquinoline, 0.58 g EDTA, 0.8 g NaOAc, 9.5 g guanidine thiocyanate, 4.6 g guanidine hydrochloride
- PGTX contains phenol, wear safety gear and gloves
- Synechocystis culture
- Chloroform/IAA
- IPA + NaOAc (in a specific amount)
- 1 vol. IPA or 3 vol. EtOH and NaOAc 30:1 or 10:1

Start

Centrifuge 5 ml of culture (OD₇₅₀ of ~ 1) for 5 minutes at maximum speed (T = 4°C). Discard supernatant. Resuspend cyanobacterial pellet in the remaining water (~1 ml). Transfer to a fresh 2 ml tube (RNase-free). Spin down 1 min at maximum speed. Discard remaining supernatant. resuspend pellet in 1 ml PGTX solution. Flash freeze and store at -80°C for later extraction, or proceed with the next step.

Extract

Heat samples at 95°C for 5 minutes in a shaking heat block. Vortex samples from time to time to ensure complete lysis.
 Place samples on ice for 5 minutes.
 Add 700 µl Chloroform/IAA. Vortex sample until it is opaque. Incubate at RT for 10 min, vortexing from time to time.
 Centrifuge samples at maximum speed for 10 minutes to separate phases. Transfer aqueous phase (600 µl) to a fresh tube (RNase-free).
 Add 1 vol. (600µl) Chloroform/IAA. Mix well by vortexing. Centrifuge 10 minutes at

maximum speed. Transfer aqueous phase (500µl) to a fresh tube (RNase-free).

Precipitation

Add 517 µl IPA + NaOAc to the samples. Mix well. Precipitate over weekend at -20°C. Also possible: 1h at -80°C or over night at -20°C.

Centrifuge precipitated sample at 4°C and maximum speed for at least 30 min.

Remove supernatant, making sure not to disrupt the RNA pellet.

Wash pellet with 70% EtOH (300µl). Centrifuge for 15 minutes, 4°C at maximum speed. Completely remove supernatant. Dry at RT for ~5 minutes. Do not overdry!

Resuspend pellet in 40 µl pure, RNase-free water. Store at -80°C.