Ki-67 Staining Protocol V.2

Sam Li¹

¹BioLegend

Sam Li
BioLegend

MATERIALS

MATERIALS

DOI:
dx.doi.org/10.17504/protocols.io.bac2iaye

External link:
<https://www.biolegend.com/protocols/ki-67-flow-cytometry-staining-protocol/4263/>

Protocol Citation: Sam Li .
Ki-67 Staining Protocol.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bac2iaye>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Dec 10, 2019

Last Modified: Dec 10, 2019

PROTOCOL integer ID:
30842

Keywords: ki-67, flow cytometry, cell division, proliferation, replicating

- 1 Prepare 70% Ethanol and chill to -20°C.
Tip: Do not freeze ethanol for long-term storage.
- 2 Prepare target cells of interest and wash 2X with PBS, centrifuging at 350xg for 5 minutes.

- 3 Discard supernatant and loosen the cell pellet by vortexing.
- 4 Add 3ml cold 70% ethanol drop by drop to the cell pellet while vortexing.
- 5 Continue vortexing for 30 seconds and then incubate at -20°C for 1 hour.
- 6 Wash 3X with BioLegend's Cell Staining Buffer (Cat. No. [420201](#)) and then resuspend the cells at the concentration of $0.5-10 \times 10^6/\text{ml}$.
- 7 Mix 100µl cell suspension with proper fluorochrome-conjugated Ki-67 antibody and incubate at room temperature in the dark for 30 minutes.
- 8 Wash 2X with BioLegend's Cell Staining Buffer and then resuspend in 0.5ml cell staining buffer for fluorescence activated cell sorting (FACS), or flow cytometric analysis.
Tip: Based on customer testing, Ki-67 staining is not recommended with our True-Nuclear™ Transcription Factor Buffer Set.