

Mar 16, 2023

Version 2

Kraus et al., 2022 FBXO7 /Park15 V.2

DOI

<https://dx.doi.org/10.17504/protocols.io.kxygx99pwg8j/v2>

Felix Kraus^{1,2}

¹Department of Cell Biology, Blavatnik Institute, Harvard Medical School, 240 Longwood Ave, Boston MA 02115, USA;

²Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD 20815, USA



Felix Kraus

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.kxygx99pwg8j/v2>

Collection Citation: Felix Kraus 2023. Kraus et al., 2022 FBXO7 /Park15 . **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.kxygx99pwg8j/v2> Version created by **Felix Kraus**

License: This is an open access collection distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this collection and it's working

Created: March 16, 2023

Last Modified: May 31, 2024

Collection Integer ID: 78871

Keywords: ASAPCRN, autophagy machinery to damaged mitochondria, ubiquitin ligase substrate receptor fbxo7, mitophagy receptor, recruitment of mitophagy receptor, ubiquitylation of mitochondrial outer membrane protein, obvious alterations in mitochondria, involvement of fbxo7, mitochondria, damaged mitochondria, mitochondrial outer membrane protein, ubiquitin ligase parkin, fbxo7 mutation, global proteomics of neurogenesis, removal of damaged mitochondria, pub puncta on mitochondria, mitochondrial clearance, dependent mitophagy, absence of fbxo7, autophagy machinery, general role for fbxo7, neurogenesis, role for fbxo7, other organelle, phosphorylation, protein, fbxo7, mitophagic flux, parkin activation, global proteomic

Funders Acknowledgements:

ASAP

Grant ID: ASAP-000282

Abstract

The protein kinase PINK1 and ubiquitin ligase Parkin promote removal of damaged mitochondria via a feed-forward mechanism involving ubiquitin (Ub) phosphorylation, Parkin activation, and ubiquitylation of mitochondrial outer membrane proteins to support recruitment of mitophagy receptors. The ubiquitin ligase substrate receptor FBXO7/PARK15 is mutated in an early-onset parkinsonian-pyramidal syndrome. Previous studies have proposed a role for FBXO7 in promoting Parkin-dependent mitophagy. Here, we systematically examine the involvement of FBXO7 in depolarization-dependent mitophagy in the well-established HeLa and induced-neurons cell systems. We find that FBXO7^{-/-} cells have no demonstrable defect in: 1) kinetics of pUb accumulation, 2) pUb puncta on mitochondria by super-resolution imaging, 3) recruitment of Parkin and autophagy machinery to damaged mitochondria, 4) mitophagic flux, and 5) mitochondrial clearance as quantified by global proteomics. Moreover, global proteomics of neurogenesis in the absence of FBXO7 reveals no obvious alterations in mitochondria or other organelles. These results argue against a general role for FBXO7 in Parkin-dependent mitophagy and point to the need for additional studies to define how FBXO7 mutations promote parkinsonian-pyramidal syndrome.

Files

 SEARCH

Protocol

NAME

Flow cytometry-based measurement of mitophagic flux

VERSION 1

CREATED BY



Felix Kraus

OPEN →

Protocol



NAME

Whole-cell proteomics and Analysis by Tandem Mass Tagging-based proteomics

VERSION 1

CREATED BY



Felix Kraus

OPEN →

Protocol

NAME

Evaluation of pUb kinetics using 3D-SIM

VERSION 1

CREATED BY



Felix Kraus

OPEN →

Protocol

NAME

Evaluation of mtKeima foci in induced neurons (iNeurons)

VERSION 1

CREATED BY



Felix Kraus

[OPEN](#) →

Protocol

NAME

Microscopy-based mtDNA turnover measurements in HeLa and iNeurons

VERSION 1

CREATED BY



Felix Kraus

[OPEN](#) →

Protocol

NAME

Microscopy-based mitochondrial morphology measurements in iNeurons

VERSION 1

CREATED BY



Felix Kraus

[OPEN](#) →

Protocol

NAME

Microscopy-based measurements of p62 recruitment in HeLa

VERSION 1

CREATED BY



Felix Kraus

[OPEN](#) →

Protocol

NAME

Microscopy-based pUb-coverage measurements of mitochondria in iNeurons

VERSION 1

CREATED BY



Felix Kraus

[OPEN](#) →

Protocol

NAME

Microscopy-based evaluation of mtKeima flux in hESC-derived Ctrl and FBXO7-/- iNeurons

VERSION 2

CREATED BY



Felix Kraus

OPEN →

Protocol

NAME

Immunocytochemical analysis

VERSION 1

CREATED BY



Felix Kraus

OPEN →