


Mar 27, 2024

🌐 Labeling of host immune cells with magnetic nanoparticles

DOI

dx.doi.org/10.17504/protocols.io.36wgq36z3lk5/v1



Eric J. Koch^{1,2}, Jamie S Foster¹

¹Department of Microbiology and Cell Science, University of Florida;

²Observatoire Oceanologique de Banyuls-sur-Mer Sorbonne Université



Jamie S Foster

University of Florida

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.36wgq36z3lk5/v1>

Protocol Citation: Eric J. Koch, Jamie S Foster 2024. Labeling of host immune cells with magnetic nanoparticles. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.36wgq36z3lk5/v1>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited



Protocol status: Working

We use this protocol and it's working

Created: September 29, 2023

Last Modified: March 27, 2024

Protocol Integer ID: 88590

Keywords: symbiosis, magnetic nanoparticles, labeling immune cells, immune cells with magnetic nanoparticle, magnetic nanoparticle, available magnetic nanoparticle, magnetic particle imaging, magnetic tweezer, hemocyte, immune cell, techniques such as magnetic tweezer, host squid, cell, cell type with phagocytic capability, innate immune cell, phagocytic capability, labeling of host, magdot, labeling technique

Funders Acknowledgements:

Moore Foundation

Grant ID: 9349

Abstract

This protocol describes a technique to label the innate immune cells of a host squid, known as hemocytes, with commercially available magnetic nanoparticles (i.e., magdots). The goal of this technique is to label the hemocytes so that they could be reintroduced into the host animal and be tracked with downstream applications, such as fluorescence and magnetic particle imaging, or manipulated with techniques such as magnetic tweezers. The labeling technique can be applied to any cell type with phagocytic capabilities.

Image Attribution

Image by Eric J. Koch. Squid hemocytes isolated from host animal, attached to slides, and counterstained with f-actin (green) and TOTO-3 nucleic acid (blue) stains. The image demonstrates the ability of host hemocytes to phagocytose magnetic nanoparticles (purple). Inset: control cells with no phagocytosed particles.



Materials

Squid Ringer's Solution

530 mM NaCl,
10 mM KCl,
25 mM MgCl₂,
10 mM CaCl₂ and
10 mM HEPES buffer,
pH 7.5

Commercially Available MagDots

Polyethylene glycol magdots - emit fluorescence at 610 nm

Note other Magdots available here:

<https://www.corequantum.com/products/magdot/>

Troubleshooting

Preparation of target immune cells - squid hemocytes

- 1 Collect sample of target immune cells. Here, hemocytes extracted from the bobtail squid *Euprymna scolopes* were used.
- 2 Using an insulin 1 cc syringe (30 gauge needle) collect hemolymph from the cephalic artery in the bobtail squid as visualized in Collins and Nyholm, 2010. Typically, from an adult animal 200 μ L can be collected with a single bleeding event. Place the hemolymph in a 1.5 mL centrifuge tube.
- 3 Resuspend the hemocytes by vigorously pipetting the hemolymph 5 to 10 times. Be sure to break up any clumps of cells.
- 4 Distribute the hemolymph into the desired number of 1.5 mL centrifuge tubes based on the experimental design (e.g., four tubes for four conditions).
- 5 Centrifuge the hemocytes at 2000 RCF, 4°C for 15 minutes to pellet the hemocytes.

Preparation of Magnetic nanoparticles (i.e., magdots)

- 6 For this protocol, polyethylene glycol-coated MagDots that fluoresce at 610 nm from Core Quantum Technologies (<https://www.corequantum.com/>) was used. Many other types of particles are available as well as other wavelengths and coatings. Note the concentration of the commercial product will vary between purchases. The goal is to have the final concentration of the magdots to be 100 μ g/mL, therefore, adjust the volumes accordingly.
- 7 Pipette 5 μ L of magdots per sample for a final concentration of 10 μ g/ 100 μ L. A typical labeling experiment would have 10 replicate hemocyte samples, therefore on average 50 μ L of magdots are used per experiment.
- 8 While the hemocytes are being pelleted, remove the total amount of magdots needed for all conditions and experimental treatments and place into a 1.5 mL centrifuge tube. Centrifuge the magdots at 10,000 RCF (or maximum speed) to pellet the magnetic particles and remove the supernatant containing the stabilization buffer that came with the particles. Note: you will likely have to leave a few μ L of buffer behind (e.g. between 10 and 20 μ L).
- 9 Resuspend the magdots in the total amount of buffer necessary for all samples. In this case, 80 μ L of Squid Ringer's solution (see materials for recipe; Nyholm et al., 2009)



was added per sample (e.g. 400 μL of Squid Ringer's would be required for five samples).

- 10 Break up the magdot pellet with either a bath sonicator for 2 min or by vigorously pipetting until the pellet is dispersed.

Co-incubation of Magnetic nanoparticles and immune cells

- 11 Immediately after the hemocytes have been pelleted, remove cell-free hemolymph leaving only 20 μL (or 20% of the incubation volume if you are not using 100 μL).
- 12 Add the 80 μL of resuspended magdots in Squid Ringer's solution to each of the prepared tubes containing 20 μL of hemocytes for a final volume of 100 μL . Note the magdot final concentration should be 100 $\mu\text{g/mL}$.
- 13 For a negative control, add 100 μL of Squid Ringer's solution with no magdots added.
- 14 Incubate the hemocytes with the magdots overnight (between 12 - 24 h) in the dark at room temperature.
- 15 Following the incubation, resuspend the hemocytes and magdots by pipetting vigorously several times.
- 16 Immediately centrifuge the samples at 500 - 1000 RCF for 15 min to pellet the cells but keeping the free, unincorporated magdots suspended.
- 17 Gently remove the supernatant containing the magdots and discard.
- 18 Rinse the now labeled hemocytes by suspending in 100 - 200 μL of Squid Ringer's solution and centrifuge for 500 - 1000 RCF for 15 min.
- 19 Repeat the rinsing process 3 to 5 times until there are no magdots visible in the supernatant.
- 20 The labeled hemocytes are now ready for visualization or downstream application.



Protocol references

Collins, A.J. and Nyholm, S.V., 2010. Obtaining hemocytes from the Hawaiian bobtail squid *Euprymna scolopes* and observing their adherence to symbiotic and non-symbiotic bacteria. *JoVE (Journal of Visualized Experiments)*, (36), p.e1714.

Nyholm, S.V., Stewart, J.J., Ruby, E.G. and McFall-Ngai, M.J., 2009. Recognition between symbiotic *Vibrio fischeri* and the haemocytes of *Euprymna scolopes*. *Environmental Microbiology*, 11(2), pp.483-493.