FEB 21, 2021

LAMP/RT-LAMP Buffer protocol

In 1 collection

Tamara Matute¹, Isaac Núñez², Fernan Federici²

¹LabTecLibres; ²SynBioUC

Laboratorio de Tecnologías Libres

Reclone.org (The Reagent Collaboration Network)

[1 more workspace](#)

Tamara Matute

ABSTRACT

This protocol describes the preparation of **reaction buffer** for homemade LAMP / RT-LAMP reactions which involves the use of home-brewed BstLF and MMLV enzymes. It also contains the instructions to prepare the enzyme **storage buffers**.

Home-brewed enzymes information:

BstLF polymerase:

expression and purification [protocol](#).

expression [plasmid sequence](#).

M-MLV reverse transcriptase:

expression and purification [protocol](#).

expression [plasmid sequence](#).

Testing

We test the buffers on SARS-CoV-2 LAMP or RT-LAMP reactions with synthetic dsDNA or ssRNA respectively.

The general performance of Homemade buffer seems appropriate and no amplification was obtained for non-template controls (NTC).

To properly formulate the buffer we tested different KCl concentrations which is known to be a critical factor in the buffer. For the home-brewed BstLF, we found that 50 mM KCl is the best concentration, displaying equal or better performance than the commercial buffer. No amplification was observed for 150 mM KCl formulation.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bsehnbb6

Protocol Citation: Tamara Matute, Isaac Núñez, Fernan Federici 2021. LAMP/RT-LAMP Buffer protocol. [protocols.io](https://dx.doi.org/10.17504/protocols.io.bsehnbb6) <https://dx.doi.org/10.17504/protocols.io.bsehnbb6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 14, 2021

Last Modified: Feb 21, 2021

PROTOCOL integer ID:
47273

Keywords: LAMP, RT-LAMP, Isothermal Amplification Buffer, IAB, Storage Buffer, ReClone, BstLF, MMLV, RNA detection, Viral detection, Home-made reactions, Enzyme storage buffer, open recipes

Figure 1. LAMP Amplification results. 10.000 copies of [plasmid synthetic sample](#) with SARS-Cov2 gen N were added as positive sample and N2 primers set [1] was used to carry the LAMP reaction. BstLF corresponds to the home-brewed enzyme and Bst2.0 warmstart to the commercial reference used. NEB IAB: NEB Isothermal Amplification Buffer. HM IAB: Homemade Isothermal Amplification Buffer. NTC: non Template control.

Figure 2. Buffer formulations effect on Threshold time (Tt) and Amplification Step size of positive LAMP reactions. Commercial NEB Isothermal Amplification Buffer (IAB) and Bst2.0 Warmstart Polymerase (Bst2.0 WS) was used to benchmark. KCl indicated concentrations correspond to 1X buffer.

[1] [Zhang, Y. et al. Enhancing colorimetric loop-mediated isothermal amplification speed and sensitivity with guanidine chloride. *Biotechniques* 69, \(2020\).](#)

IMAGE ATTRIBUTION

Tamara Matute & Isaac Núñez. GNU General Public License v3.0.

MATERIALS

- pH-meter
- Magnetic Stirrer
- Precision Scale
- 0.22 uM filters
- Syringe
- Autoclave
- Shott Bottles
- 15 and 50 ml conical tubes

BstLF and MMLV storage buffer

- 1 Enzyme storage buffer was made based on [NEB indicated Enzymes Storage Conditions](#).

Component	10X Storage Buffer	1x Storage Buffer
Tris-HCl (pH 7.1 @ 25°C)	100 mM	10 mM
KCl	500 mM	50 mM
EDTA	1 mM	0.1 mM
Triton® X-100	1%	0.1%
Glycerol	-	50%
DTT	-	1 mM

Note

DTT should be added as fresh as possible. Add DTT and Glycerol at the moment of assembling the 1X Buffer. Freeze and thaw cycles are not recommended for DTT solutions, always use a fresh aliquot.

- 2 To prepare 10 mL of **10x Storage Buffer** add:
- 6 mL molecular grade H₂O
 - 1 mL 1M Tris-HCl pH 7.1
 - 2 mL KCl 2.5M
 - 20 µL EDTA 500 mM
 - 100 µL Triton® X-100
 - Make up the volume to 10 mL with H₂O (~ 880 µL)
 - Make aliquots and store at -20 °C

Note

In the "**Stocks Solutions**" section you can find the preparation of each solution.

- 3 To prepare 1.5 mL of **1x Storage Buffer** add:
- 599 µL molecular grade H₂O
 - 150 µL 10x Storage Buffer
 - 1.5 µL DTT 1M
 - 500 µL Glycerol 100%
 - Mix properly (by vortex), use immediately or store at -20 °C

LAMP and RT-LAMP reaction buffer

- 4 This (RT-)LAMP assay buffer was made based on [NEB 1X Isothermal Amplification Buffer Pack](#).

10X Isothermal Amplification Buffer is as follow:

Component	10X Buffer concentration	1X Reaction concentration
Tris-HCl (pH 8.8.@25°C)	200 mM	20 mM
(NH ₄) ₂ SO ₄	100 mM	10 mM
KCl	500 mM	50 mM
MgSO ₄	20 mM	2 mM
Tween® 20	1%	0.1%

- 5 Prepare the following **stock solutions**:

Component	Concentration
Tris-HCl (pH 8.8.@25°C)	1M
(NH ₄) ₂ SO ₄	1M
KCl	2.5M
MgSO ₄	1M
Tween® 20	100%

Note

In the "**Stocks Solutions**" section you can find the preparation of each solution.

6 To prepare 50 mL of 10X Isothermal Amplification Buffer add:

- 20 mL Molecular H₂O
- 10 mL 1M Tris-HCl (pH 8.8 @ 25°C)
- 5 mL 1M (NH₄)₂SO₄
- 10 mL 2.5M KCl
- 1 mL 1M MgSO₄
- 0.5 mL Tween® 20
- Adjust pH 8.8 with NaOH 1 M (~ 560 µL)
- Make up the volume to 50 mL with H₂O (~ 3 mL)
- Filter-sterilize the solution with 0.22 µm filter.

Note

It shouldn't be autoclaved.

The autoclave is not recommended for Tween® 20, (NH₄)₂SO₄ and MgSO₄ solutions.

- Make aliquots and store at -20 °C.

pH variation dynamics (@25 °C) along buffer elaboration.

Stock solutions

7 Potassium chloride (KCl)

To prepare 50 ml of a 1 M solution, weigh 9.3188 g of KCl powder and completely dissolve it in 40 mL of H₂O. Make up the volume to 50 mL with H₂O (it can be done in a 50 ml conical tube). Filter-sterilize the solution with a 0.22 µm filter and store at room temperature.

8 Ammonium sulfate ((NH₄)₂SO₄)

To prepare 50 ml of a 1 M solution, weigh 6.6070 g of (NH₄)₂SO₄ powder and completely dissolve it in 40 mL of H₂O. Make up the volume to 50 mL with H₂O (it can be done in a 50 ml conical tube). Filter-sterilize the solution with a 0.22 µm filter and store at room temperature.

9 Magnesium sulfate (MgSO₄)

To prepare 50 ml of a 1 M solution, weigh 12.3240 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ powder and completely dissolve it in 40 mL of H_2O . Make up the volume to 50 mL with H_2O (it can be done in a 50 ml conical tube). Filter-sterilize the solution with a 0.22 μm filter and store at room temperature.

Note

It is possible to use MgSO_4 instead of heptahydrated but it is more hygroscopic and would need desiccating before weighing.

10 Tris-HCl

To prepare 250 ml of a 1 M solution:

- Weigh 30.2850 g of Trizma Base powder and completely dissolve it in 200 mL of H_2O . It should be done in a magnetic stirrer.
- Adjust to pH 8.8 or 7.1 with concentrated HCl (2.5 M). It should be done in a properly calibrated pH-meter, with continuous stir and giving enough time to let the pH to equilibrate. It is important to check the temperature with a thermometer and use a heater if necessary to reach a temperature solution of 25°C the whole time.
- Transfer the solution to a graduated cylinder and complete the volume to 250 ml with H_2O .
- Filter-sterilize the solution with a 0.22 μm filter
- Store at room temperature or 4°C up to six months.

Note

Temperature has a significant effect on the pH of Tris buffers. It decreases ~0.028 pH units per each °C increase.

11 Hydrochloric acid (HCl)

To prepare 50 ml of 2.5 M solution:

- Weigh 4.55 g HCl and completely dissolve it in 40 mL of H_2O , or add 10.25 mL HCl 37% ($d = 1.19$) solution to 40 ml H_2O .
- Make up the volume to 50 ml with H_2O .

12 EDTA (ethylenediaminetetraacetic acid)

To prepare 500 ml of a 500 mM solution:

- Weigh 93.06 g of EDTA disodium salt dihydrate powder ($\text{MW} = 372.24 \text{ g/mol}$) and dissolve it in 400 mL of H_2O . At this concentration it shouldn't dissolve completely until you up the pH.
- Adjust to pH 8.0 or 7.1 with concentrated NaOH. It would be needed ~25 μL of 10M NaOH. It should be added slowly in a magnetic stirrer letting the EDTA to dissolve and the pH to equilibrate.
- Transfer the solution to a graduated cylinder and complete the volume to 500 ml with H_2O .
- Filter-sterilize the solution with a 0.22 μm filter
- Store at room temperature or 4°C.

13 DTT (dithiothreitol)

To prepare 10 ml of 1 M solution, weigh 1.54 g of DTT powder and dissolve it completely in 80 mL of H_2O . Make up the volume to 10 mL with H_2O (it can be done in a 15 ml conical tube). Filter-sterilize the solution with a 0.22 μm filter, make 100 μL - 1 ml aliquots and store at -20°C.

14 Glycerol

Make an aliquot and sterilize by autoclave.

15 Filters:

To sterilize the solutions and the assembled buffer 0.22 µm syringe filters of MCE or Cellulose acetate membranes could be used.

Specifically we used:

BioFil

MCE membrane

0.22 µm

EDLAB

Cellulose acetate

0.22 µm