NOV 28, 2018

Limiting Dilution & Clonal Expansion

Synthego¹

¹Synthego

Synthego

Brittany Enzmann

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.srqed5w

External link:

<https://www.synthego.com/resources/Limiting-Dilution-&-Clonal-Expansion-Protocol>

Protocol Citation: Synthego 2018. Limiting Dilution & Clonal Expansion.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.srqed5w>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol in our group and it is working.

Created: Aug 20, 2018

Last Modified: Nov 28, 2018

PROTOCOL integer ID:

14864

Keywords: CRISPR-Cas9, CRISPR, isolation of single cells, limiting dilution, clonal expansion

ABSTRACT

CRISPR-Cas9 can be used to generate edited cell pools for a variety of applications. These pools contain a heterogeneous mix of cells, some unedited and some with different genetic alterations at the targeted locus. In order to produce a cell line with a particular edit or knockout, clonal expansion of individual cells is necessary. This protocol describes how to isolate single cells from a CRISPR-edited pool, and expand and genotype the subsequent clonal populations.

The isolation of single cells from a knockout cell pool can be accomplished through limiting dilution. While there are more sophisticated means to accomplish single cell plating (e.g., flow cytometry), limiting dilution is a technique that can be carried out with standard pipetting tools. The obvious drawback to this technique is that it relies on statistical probability (based on a Poisson distribution) to deposit a single cell per well. A typical concentration used for limiting dilution is 0.5-1 cells per 100 μ l aliquot because it maximizes the probability of obtaining a single cell per well while minimizing the probability of depositing multiple cells per well. At this concentration, one can expect about a third of the wells to contain a single cell and most wells to be empty. It is much less difficult to exclude wells without cells than to sort out heterogeneous populations arising from multiple cells.

In this protocol, cells from each edited population are diluted to 0.5-1 cells per 100 μ l and plated on at least two 96-well plates (100 μ l/well). However, if clonally expanding cells for the first time we recommend plating a higher number of cells (e.g., 4 x 96-well plates per condition).

To control for potential health/growth defects caused by the editing, we recommend including mock and/or negative control for single cell dilution, although these do not require multiple 96-well plates. The *RELA* positive control, however, does not need to be plated out for clonal expansion as it serves mainly as a positive control for transfection and editing efficiency.

ATTACHMENTS

[Limiting Dilution and Clonal Expansion.pdf](#)

GUIDELINES

Important Considerations

The ease at which a clonal cell line can be developed depends on the cell type. It is critical that the cells are healthy and have reached their respective doubling rate before limiting dilution is attempted. For most cell types, waiting 6 days after transfection is typically enough time for the cells to stabilize. Ideally, limiting dilution and clonal expansion should be conducted after 1 passage. Clonal expansion from a single cell can take anywhere from 2-8 weeks depending on the cell type. It is important to continually monitor the health and growth of clones during expansion.

It is important to consider that edits do not necessarily equate to knockouts. The insertion or deletion (indel) of a number of nucleotides that is not divisible by three will cause a frameshift mutation in the target gene. The resultant protein will most likely be nonfunctional.

Pools with the highest indel frequencies generate clonal cell lines with desired edits most efficiently. Synthego Knockout Cell Pools guarantee a genotypic and protein knockout of at least 50%.

Synthego's [Inference of CRISPR Edits \(ICE\) tool](#) can accurately characterize edited cell pools and clones. Learn more about ICE [here](#).

The protocol workflow is as follows:

1. Limiting Dilution (1 Day)
2. Clonal Expansion (2-8 weeks)
3. Genotyping Clones
4. Storage & Additional Analyses

Additional Information

For an up-to-date list of all Synthego Protocols and other resources, please visit synthego.com/resources

For technical assistance, contact our Scientific Support Team:

Ph: 888.611.6883

Email: support@synthego.com

MATERIALS

MATERIALS

- ✕ 1.5 ml microcentrifuge tubes **Contributed by users**
- ✕ 10 ml sterile reagent reservoir **Contributed by users**
- ✕ Multichannel pipette or similar **Contributed by users**
- ✕ 96-well cell culture plates (up to 25) **Contributed by users**
- ✕ Normal growth medium **Contributed by users**
- ✕ Phosphate buffered saline without calcium and magnesium (PBS) **Contributed by users**
- ✕ TrypLE™ cell dissociation reagent **Life Technologies Catalog #12605010**
- ✕ Hemocytometer or automated cytometer **Contributed by users**
- ✕ Trypan blue **Contributed by users**
- ✕ Cell strainer (optional) **Contributed by users**

SAFETY WARNINGS

Please refer to the SDS (Safety Data Sheet) for safety warnings and hazard information.

Limiting Dilution

- 1 Warm medium at 37°C in a water bath.
37 °C water bath
- 2 Transfer 24-well plate containing transfected cells from the humidified CO₂ incubator to a biological safety cabinet.
- 3 Aspirate medium from cells across all conditions.

- 4 Carefully wash cells with PBS.
- 5 Aspirate PBS.
- 6 Add 250 µl of TrypLE™ (or other appropriate cell dissociation reagent) to cells (adjust volumes of TrypLE™ to the size of the cell culture vessel if cells have been passaged).

250 µL TrypLE
- 7 Incubate the cells for 5 minutes at 37°C and 5% CO₂ or until the cells detach.

37 °C 5% CO2 Incubation

00:05:00 5% CO2 Incubation
- 8 Transfer well plate back to the biological safety cabinet.
- 9 To neutralize dissociation, add an equal volume of medium (250 µl) to that of the TrypLE™ to each of the wells.

250 µL medium
- 10 Pipet 3-4 times (more if needed) with 1 ml pipette to break up cell clumps. Cells can also be passed through a cell strainer.
- 11 Transfer the cells from each condition to a separate sterile 1.5 ml microcentrifuge tube.
- 12 Count the cells from each condition using a hemocytometer or automated cytometer.

- 13 Calculate the concentration of cells (cells/ml) in the cell suspension.
- 14 Determine the number of 96-well plates for each pool. It is recommended to plate one or two 96-well plates for each edited pool to increase the likelihood that at least one successful knockout clone is generated.
- 15 Dilute cells in a series of 10-fold dilutions to a final concentration of 0.5-1 cells/100 μ l of medium (this will give around 1 cell/well in most of the wells in a 96-well plate) or 120 cells in 12 ml of normal growth medium. Scale according to the number of plates desired. For example, if 5 x 96-well plates are desired for a particular condition, dilute 600 cells in 60 ml of normal growth medium.

Note

The concentration of cells can be adjusted (0.5-5 cells/100 μ l).

- 16 Transfer the diluted cell suspension to a large sterile reservoir.
- 17 Dispense 100 μ l/well of the diluted cell suspension to each well using a multichannel pipette.
- 18 Repeat steps 13 to 17 for each edited population selected for clonal expansion.

⇒ go to step #13 Repeat step 13 to 17

- 19 Transfer plates to a humidified incubator at 37°C, 5% CO₂. Change medium as needed.

🔥 37 °C 5% CO2 Incubator

- 20 Expand & genotype clones (see below).

Clonal Expansion

- 21 Visually screen plates using a microscope for established single colonies for about two weeks. If a clone imager is available, we recommend imaging each of the 96-well plates 2-4 hours 02:00:00 - 04:00:00 after plating the single cell dilution (before the cell can divide). The cells should be imaged 5-7 days thereafter. This practice enables one to track cell growth over time and confirm that the final clones originated from a single cell.

Mark establishing colonies by circling the well on the lid and/or record in a spreadsheet. Note that different cells (or possibly genetically modified cells) may grow at a different pace. When colonies reach 70% confluence, transfer them to a 24-well plate. During transfer, take out a portion of cells from each colony for genotyping (see below). By continuing to expand only colonies with confirmed edits, one can reduce the burden of culturing and future screening assays. Maintain a few colonies that grow from the Mock or Negative Control conditions to be used as an isogenic wild type comparison to the knockout cell line.

Genotyping Clones

- 22 To genotype your clones, isolate genomic DNA, PCR-amplify the edited region, and sequence the amplicons via Sanger sequencing. Synthego's [Inference of CRISPR edits \(ICE\)](#) tool can be used to analyze the sequence data.

Note

A clone can be reported as wild type (unedited), containing a homozygous edit, or containing a heterozygous edit. A homozygous edit is when both alleles of a chromosomal pair have the same mutation. These colonies should be saved and marked as potential knockouts. A heterozygous edit is when one allele has a mutation and the other does not (wild type), or when both alleles have different mutations (i.e., compound heterozygous). The heterozygous edit can be easily identified by the overlapping peaks in the chromatogram centered around the region targeted by the guide. Double-stranded breaks repaired by the NHEJ pathway often result in heterozygous edits. This is especially true when editing hyperploid cell lines. Because compound heterozygous mutations may inactivate genes, such colonies should be marked as potential knockouts.

Note

Whether the edits are of homo- or heterozygous nature, it is recommended to select colonies containing indels that give rise to frameshift mutations, ideally those that generate a premature stop codon. Indels that maintain the gene's reading frame or alter the coding of only a few amino acids towards the 3' end of the exon, may not lead to the loss of the protein's function.

Storage & Additional Analyses

- 23** After identifying putative knockouts by genotype, the cells should be split and expanded for cryopreservation while performing further analysis to functionally validate the knockout. Functional validation of a knockout can be achieved using immunoblot, ELISA or flow cytometry analysis or an appropriate activity assay when available.