VERSION 2

NOV 04, 2017

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.kk3cuyn

Protocol Citation: Emilie Viennois 2017. Lithium chloride purification, a rapid and efficient technique to purify RNA from DSS-treated tissues. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.kk3cuyn>

MANUSCRIPT CITATION: Dextran sodium sulfate inhibits the activities of both polymerase and reverse transcriptase: lithium chloride purification, a rapid and efficient technique to purify RNA. *Viennois et al. BMC Research Notes* 2013, **6**:360. doi:10.1186/1756-0500-6-360

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Lithium chloride purification, a rapid and efficient technique to purify RNA from DSS-treated tissues V.2

Emilie Viennois¹

¹Georgia State University

Emilie Viennois
 Georgia State University

ABSTRACT

Dextran sodium sulfate (DSS) is commonly used in mouse studies to induce a very reproducible colitis that effectively mimics the clinical and histological features of human inflammatory bowel disease (IBD) patients, especially ulcerative colitis. However, the mechanisms of action of DSS remain poorly understood, and observations by our laboratory and other groups indicate that DSS contamination of colonic tissues from DSS-treated mice potentially inhibits the quantitative reverse-transcription polymerase chain reaction (qRT-PCR) amplification of mRNA. A prior study used poly-A-mediated mRNA purification to remove DSS from RNA extracts, but we herein report a second efficient and cost-effective approach to counteract this inhibition, using lithium chloride precipitation to entirely remove DSS from RNAs. We also explored how DSS interferes with qRT-PCR process, and we report for the first time that DSS can alter the binding of reverse transcriptase to previously primed RNA and specifically inhibits the enzymatic activities of reverse transcriptase and Taq polymerase in vitro. This likely explains why DSS-treated colonic RNA is not suitable to qRT-PCR amplification without a previous purification step. In summary, we provide a simple method to remove DSS from colonic RNAs, and we demonstrate for the first time that DSS can inhibit the activities of both polymerase and reverse transcriptase. In order to reliably analyze gene expression in the colonic mucosa of DSS-treated mice, the efficiency rate of qRT-PCR must be the same between all the different experimental groups, including the water-treated control group, suggesting that whatever the duration and the percentage of the DSS treatment, RNAs must be purified.

Created: Nov 04, 2017

Last Modified: Mar 17, 2018

PROTOCOL integer ID:
8571

Keywords: DSS, DNA-polymerase, Reverse transcriptase, Lithium chloride, RNA

ATTACHMENTS

[Viennois E 2013 DSS and LiCl.pdf](#)

GUIDELINES

DSS treated mice

All studies were performed in accordance with the Institutional Animal Care and Use Committee at Georgia State University (Atlanta, GA). All procedures were approved and are registered in the protocol IACUC ID: A11025, approval date 8/30/2011 to 8/30/2014. Strains, ages, and the number of animals follow the established protocol. The DSS treatment on mice were carried out in C57BL/6 mice (8 wk, 18–22 g) obtained from Jackson Laboratories (Bar Harbor, ME). Mice were group housed under a controlled temperature (25°C) and photoperiod (12:12-h light–dark cycle) and allowed unrestricted access to standard mouse chow and tap water. DSS [40,000 Da, 3% (wt/vol), ICN Biochemicals, Aurora, OH] was diluted at 3% in drinking water. After 7 days under DSS treatment, the mice were sacrificed by CO₂ euthanasia. A small piece (50 mg) of proximal colon was taken for RNA extraction.

RNA extraction

Total RNA was isolated from colonic tissues using TRIzol (Life Technologies, Carlsbad, CA) according to the manufacturer's instructions. Where indicated, RNA was purified via precipitation with lithium chloride. The RNA integrity was assessed by 2% agarose gel electrophoresis.

cDNA synthesis and qPCR

cDNA were synthesized using the Maxima First-Strand cDNA Synthesis Kit (Thermo Scientific, Waltham, MA) according to the manufacturer's instructions. Expression of the total RNA was quantified by qPCR using Maxima SYBR Green/ROX qPCR Master Mix (Thermo Scientific) in a Realplex Thermal Cycler (Eppendorf, Hauppauge, NY). The qPCR primer sequences for 36B4 were 36B4-F: TCCAGGCTTTGGGCATCA and 36B4-R: CTTTATCA GCTGCACATCACTCAGA.

Polymerase assay

The polymerase enzymatic activity was measured using the EvaEZ™ Fluorometric Polymerase Activity Assay kit (Biotium, Hayward, CA) according to the manufacturer's instructions. Briefly, 0.01 units (10 mU) of recombinant Taq DNA polymerase (Thermo Scientific) were incubated with water or DSS (0.01 to 1 g/L). Dextran was used as a control. The enzymatic activity was quantified by fluorescence using the Realplex Thermal Cycler (Eppendorf). The fluorescence was read every 1 min for 60 min during the elongation step at 72°C.

Reverse transcriptase assay

The reverse transcriptase enzymatic activity was measured using the EnzChek

Reverse Transcriptase Assay kit (Life Technologies, Carlsbad, CA) according to the manufacturer's instructions. Briefly, 0.4 units of the reverse transcriptase, M-MuLV RT (Thermo Scientific) were incubated at 25°C with water or DSS (0.00005 g/L or 0.0005 g/L) in presence of a standardized polyA RNA template, oligodT and polymerization buffer. The enzymatic activity was quantified by fluorescence using a Synergy 2 Multi-Mode Microplate Reader (Biotek).

Surface plasmon resonance (SPR)

For SPR experiments, gold sensor chips were used (Biosensing Instrument, Tempe, AZ, USA). Briefly, the principle of this technique is the following: A first molecule is coupled to the gold sensor surface. The solution containing the second molecule (the analyte) then is flowed over the surface. This creates a mass change on the sensor surface as the two molecules interact, which is detected in real time as a deflection of the resonance angle in mDeg. In that specific experiment, the gold chip was cleaned and treated as previously described [22-24]. After placing a chip into the BI-2000 SPR (Biosensing Instrument) machine each gold biosensor chip covered with carboxydextran was activated using a mixture of 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide/N-hydroxysuccinimide (EDC/NHS) to form amide linkages between purified protein and the chipbound carboxydextran. Two successive injections of 2,400 units of M-MuLV Reverse Transcriptase (thermo Scientific) each were performed. The reverse transcriptase was previously purified using Slide-A-Lyser® Mini dialysis devices (ThermoScientific) and suspended in PBS. After coating the chip with reverse transcriptase, RNA, previously primed using random primers (Thermo Scientific) incubated at 42°C, or primed RNA incubated with DSS in different concentrations (0.05, 0.5, 5 g/L) were passed over the chip twice. A two-step interaction curve was obtained. The first step involved adsorption of primed RNA to the maximal level. In the second step, when the flow of primed RNA concentration returned to zero, nonspecific adsorbed primed RNA were released with the running buffer. The deviation of the resonance angle thus decreased to a plateau located at a level above the initial baseline. We assessed the laser deflection as directly correlated to the binding level. We thus used the laser deviation angle as the optimal parameter for the binding affinity. All comparisons between the different solutions of primed RNA and primed RNA with different concentrations of DSS were performed as a measure of the laser deviation in mDegrees (mDeg).

Statistical analysis

Values were expressed as means $\pm$ standard error of mean (SEM). Statistical analysis was performed using an unpaired two-tailed t-test by GraphPad Prism 5 software. $p < 0.05$ was considered statistically significant.

MATERIALS

MATERIALS

DSS Contributed by users

Lithium chloride purification

- 1 In order to be purified from all polysaccharides including DSS, a purification using lithium chloride is performed. First, incubate the RNA with 0.1 volume of 8 M LiCl diluted in RNase-free water on ice for 2 h. (1/2)
02:00:00 Incubation on ice
- 2 Centrifuge at 14,000 g for 30 min at 4°C. (1/2)
00:30:00 Centrifugation
- 3 Pour the supernatants out and dissolve the pellet of RNA in 200 µL of RNase-free water. (1/2)
200 µL RNase-free water
- 4 Incubate the RNA with 0.1 volume of 8 M LiCl diluted in RNase-free water on ice for 2 h. (2/2)
02:00:00 Incubation
- 5 Centrifuge at 14,000 g for 30 min at 4°C. (2/2)
00:30:00 Centrifugation
- 6 Pour the supernatants out and dissolve the pellet of RNA in 200 µL of RNase-free water. (2/2)
200 µL RNase-free water
- 7 Precipitate the RNA at – 20°C for 30 min, in 0.1 volume of 3 M sodium acetate (pH 5.2) and 2 volumes of 100% absolute ethanol.
00:30:00 Precipitation

8 Centrifuge the RNA at 14,000 g for 30 min at 4°C.

00:30:00 Centrifugation

9 Pur the supernatants out and wash the pellets with 100 µL of 70% ethanol.

100 µL 70% ethanol

10 Centrifuge at 14,000 g for 10 min at 4°C.

00:10:00 Centrifugation

11 Remove the supernatants and dissolve the RNA in 20– 50 µl of RNase-free water.

50 µL RNase-free water