SEP 24, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.4wegxbe

Protocol Citation: Eftychis Frangedakis, Susana Sauret-Gueto, Anthony West, Nicola Patron, Marta Tomaselli, Marius Rebmann, Jim Haseloff 2019. Loop L1 (odd level) type IIS cloning - pCk-ye vectors. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.4wegxbe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jun 28, 2019

Last Modified: Sep 24, 2019

PROTOCOL integer ID:
 25254

🌐 Loop L1 (odd level) type IIS cloning - pCk-ye vectors

Eftychis Frangedakis¹, Susana Sauret-Gueto², Anthony West², Nicola Patron², Marta Tomaselli³, Marius Rebmann², Jim Haseloff²

¹University of Cambridge;

²Plant Sciences, University of Cambridge, OpenPlant;

³University of Cambridge, Open Plant

OpenPlant Project

Eftychis Frangedakis
 University of Cambridge, Plant Sciences

ABSTRACT

Protocol based on:

[Pollak B, Cerda A, Delmans M, et al \(2019\) Loop assembly: a simple and open system for recursive fabrication of DNA circuits. New Phytol 222:628–640](#)

<https://doi.org/10.1111/nph.15625>

MATERIALS

MATERIALS

✂ BsaI - 5,000 units **New England Biolabs Catalog #R0535L**

✂ T4 DNA Ligase - 20,000 units **New England Biolabs Catalog #M0202S**

✂ Sterile water **Contributed by users**

✂ BSA, molecular biology grade, 20 mg/ml **New England Biolabs Catalog #B9000S**

✂ 10X NEB T4 DNA ligase buffer **New England Biolabs**

1 Determine the concentrations of the DNA parts by spectrophotometry (Nanodrop).

- 2 Prepare aliquots for DNA parts to be assembled at a concentration of 15 nM and of the pCk vector at a concentration of 7.5 nM.

To calculate the concentration needed for each part to assemble (not the backbone) in ng/ μ L, divide the length of the plasmid where they are cloned by 100. For the backbone, divide the plasmid length by 200. In this way you can prepare aliquots of your parts and backbones and add 1 μ L of each one to the plasmid mix.

- 3 . Prepare Loop assembly Level 1 reaction master mix (MM) according to Table, if four or less number of parts are assembled into a pCk-pe vector.

Com pone nts	Volu me (μ L)
Steril e water	3
10x T4 ligase buffer (NEB)	1
1 mg/m L bovin e seru m album in (NEB)	0.5
T4 DNA ligase at 400 U/ μ L (NEB)	0.25
10 U/ μ L BsaI (NEB)	0.25

Final volum e	5
---------------------	---

- 4 Prepare plasmids mix for each reaction, by adding in a 0.2 mL tube, 1 μ L of each DNA part aliquot (see step 2), 1 μ L of the pCk-pe vector and sterile water up to 5 μ L. Mix well.
- 5 Add 5 μ L of master mix to the 5 μ L of plasmids mix, to a final volume of 10 μ L. Mix well. If more than 4 DNA parts are to be assembled into a pCk-pe vector, reduce the water volume in the MM by 1 μ L for each extra 1 μ L of DNA part added in the plasmids mix.
- 6 Place samples in a thermocycler and use the following program:
Assembly: 26 cycles of 37 °C for 3 min and 16 °C for 4 min.
Termination and enzyme denaturation: 50 °C for 5 min and 80 °C for 10 min.
- 7 Transform 20 μ L of chemically competent *E. coli* cells (transformation efficiency of 1×10^7 transformants/ μ g plasmid DNA) using 2 μ L of the Loop assembly reaction and then plate on LB agar plates containing 50 μ g/mL kanamycin.
- 8 Incubate overnight at 37 °C.
- 9 Colonies with white color are likely to contain the vector with the insert while yellow color colonies will contain the empty vector.
- 10 Confirm the presence of the correct insert with Sanger sequencing using the primers pC_F and pC_R
- 11