VERSION 2

OCT 21, 2019

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.8izhuf6

External link:

<https://www.diacomp.org/shared/document.aspx?id=19&docType=Protocol>

Protocol Citation: Frank Brosius 2019. Low-Dose Streptozotocin Induction Protocol (mouse).

protocols.io

<https://dx.doi.org/10.17504/protocols.io.8izhuf6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 21, 2019

Last Modified: Oct 21, 2019

Low-Dose Streptozotocin Induction Protocol (mouse) V.2

Frank Brosius¹

¹University of Arizona

Diabetic Complications Consortium

Tech. support email: rmcindoe@augusta.edu

Lili Liang

ABSTRACT

Summary:

This protocol is used by DiaComp members to induce diabetes in a number of the animal models developed by the consortium. STZ is toxic to the insulinproducing beta cells of the pancreas and used to induce a diabetes similar to a type I diabetic ([Reference](#)). Some reports suggest cellular toxicity outside of the pancreas. STZ also exhibits broad spectrum antibacterial properties and alters the gut microbiota. Please ensure that appropriate controls are included in all studies and complementary models considered (e.g. the **Ins2^{Akita}** mouse).

Diabetic Complications:

Cardiovascular

Nephropathy

Neuropathy

Retinopathy

Uropathy

Wound-Healing

Pediatric Endocrinology

MATERIALS

Keywords: Low-Dose Streptozotocin, Diabete, Cardiovascular, Nephropathy, Neuropathy, Retinopathy, Uropathy, Wound-Healing, Pediatric Endocrinology

MATERIALS

⊗ STZ-Na-Citrate Solution **Contributed by users** **Catalog #See Below**

⊗ TB or insulin syringes w/ needle attached (26-28 ½ gauge) **Contributed by users**

⊗ Isoflurane drop jar **Contributed by users**

⊗ Isoflurane **Contributed by users**

Reagent Preparation:

Na-Citrate Buffer:

Reagents and Materials

Reagent/Material	Quantity Required	Vendor	Stock Number
Na Citrate (enzyme grade)	1.47g for 50ml	Fisher Scientific	# BP 327-1
Deionized water (ddH ₂ O)	50ml		
pH meter			

Procedure

- Dissolve 1.47 g of Na Citrate in 50 ml ddH₂O
- Test pH with pH meter, adjust buffer to 4.5 pH with monohydrate Na Citrate solution if necessary.
- Buffer should be made fresh with every group of injections
- Place appropriate amount of buffer into a sterile conical tube.

Streptozotocin (STZ):

Reagents and Materials

Reagent/Material	Quantity Required	Vendor	Stock Number
Streptozotocin		Sigma	# S-0130
Eppendorf tube			
Aluminum foil			

Procedure

Concentration: 7.5 mg/ml

- STZ should be stored at –20°C to avoid dessication.
- Weigh the appropriate amount of STZ so your final concentration in the Na-Citrate buffer will be 7.5 mg/ml and place this into an Eppendorf tube; cover with aluminum foil (light sensitive).

STZ Na-Citrate Solution:

Reagents and Materials

Reagent/Material	Quantity Required	Vendor	Stock Number
Na-Citrate Buffer from above			
STZ weighed from above			
3 ml 23 gauge syringe/needle			
Empty sterile vial			

Procedure

Concentration: 7.5 mg/ml Dosage: 50 mg/kg mouse

- Do not mix STZ into buffer until ready for injections; the drug degenerates within 15-20 minutes in solution.
- Pour contents of Eppendorf tube (STZ) into the conical (buffer).
- Mix well
- Aspirate the solution using a 3 ml 23 gauge syringe/needle (may need to repeat this several times depending on amount of solution)
- Inject contents into the empty sterile vial
- Vial contains solution for injecting the mice

BEFORE START INSTRUCTIONS

IMPORTANT: Mice should fast for four (4) hours prior to STZ induction. *The consortium has agreed that a proper fast for the mice is 4-6 hours. Some of the other institutions may fast for 6 hours; we have chosen 4 hours as a proper fasting time.* The STZ-Na Citrate buffer solution should only be prepared **immediately** before injection as the drug degrades after 15-20 minutes in the Na-Citrate buffer.

- 1 Prepare the buffers and solutions as described below. Please note that the STZNa-Citrate solution should be prepared immediately prior to injection so as to avoid degradation of the STZ.
- 2 Mice should be fasted prior to injection; four hours is usually sufficient.
- 3 Place one mouse in the Isoflurane drop jar, according to your local IACUC anesthesia standards.
- 4 Remove mouse when breathing has slowed and animal is anesthetized

- 5 Inject appropriate amount of the STZ solution IP so the final dosage is **50 mg/kg mouse**
- 6 Allow mouse to awaken and place back in cage
- 7 Repeat procedure for each animal; keep in mind that Isoflurane will need to be reloaded after every third or fourth animal for proper anesthesia.
- 8 Each mouse should be given one injection for 5 days consecutively.
- 9 Supply mice with 10% sucrose water, if necessary, to avoid sudden hypoglycemia post-injection.
- 10 **Mice should be tested for sufficient levels of hyperglycemia at 4 weeks postinjection.**
If any of the experimental animals are not sufficiently diabetic, the low dose STZ protocol should be re-administered to those mice only at week 7 post injection to obtain proper levels of hyperglycemia.