AUG 04, 2020

Low-input DNA extractions in 96-well plates

Forked from [Honeybee DNA extractions in 96-well plates](#)

Reuben McKay

Tom Harrop¹, Vercoe¹,

Ciarán Cuddy¹

¹University of Otago

Tom Harrop

MATERIALS

MATERIALS

3.2 mm stainless steel beads, RNase free Contributed by users Catalog #NEXSSB32-RNA

96 Well 0.8mL Plate (Bulk) Thermo Fisher Catalog #AB0859

ZymoBIOMICS Lysis Solution Zymo Research Catalog #D4300-1-150

Quick-DNA Magbead Plus Kit Zymo Research Catalog #D4082

RNase A solution in 10 mM Tris-HCl, pH 7.5, 15 mM NaCl at a concentration of 10 mg/mL:

250 mg RNase A
250 µL 1 M Tris-HCl pH 7.5
75 µL 5 M NaCl
Make up to 25 mL

Proteinase K solution in 50 mM Tris, pH 8, 3 mM CaCl₂, 50% Glycerol at a concentration of 20 mg/mL:

100 mg Proteinase K
250 µL 1 M Tris-HCl pH 8
6 µL 2.5 M CaCl₂
Make up to 5 mL

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.birqkd5w

Protocol Citation: Tom Harrop, Reuben McKay Vercoe, Ciarán Cuddy 2020. Low-input DNA extractions in 96-well plates. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.birqkd5w>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jul 19, 2020

Last Modified: Aug 04, 2020

PROTOCOL integer ID:
39440

Prepare lysate for extraction

1 Prepare lysate for extraction

1.1 Prepare 100 µL lysis buffer per sample.

Lysis solution	99 µL
RNase solution (10 mg/mL)	1 µL
Total	100 µL

The Zymo lysis solution can be bought separately.

This protocol also works with the Zymo Solid Tissue Buffer II that is supplied with the Quick-DNA Magbead Plus kit. Solid Tissue Buffer II comes as a 2x concentrate and has to be diluted with nuclease-free water.

1.2 Dissect e.g. 1 *M. hyperodae* pupa for a single sample.

Place up to 88 individuals in a 1000 µL, round well deepwell plate

Add 2 3.2 mm stainless steel ball bearings and 100 µL of lysis buffer with RNase to each well.

1.3 Homogenise the tissue for **90 seconds at 1200 RPM** using a plate-compatible tissue homogenizer, e.g. SPEX SamplePrep 2010 Geno/Grinder.

00:01:30

1.4 Seal the plate and mix on the plate shaker. Make sure the paste is resuspended.

Incubate at 37°C for 30 minutes.

00:30:00

37 °C

1.5 Add 5 µL Proteinase K solution (20 mg/mL) and mix on the plate shaker.

Incubate at 55°C for 120 minutes, shaking the plate for one minute every 30 minutes.

02:00:00

55 °C

1.6 Spin the plate for 10 minutes at 1,000 RPM.

1000 rpm, 15°C,
00:10:00

1.7 Transfer 100 µL of lysate to a 0.8 mL deepwell plate.

DNA extraction

2 Set up the reagents from the ZYMO *Quick*-DNA Magbead Plus Kit

2.1 Add the following reagents to the 7-position ReservoirRack

Reagent	Reservoir volume (mL)	8	16	24	32	40	48	56	64	72	80	88
1: 300 µL MBB : 5 µl beads	30	264 5	508 5	752 5	996 5	124 05	148 45	172 85	197 25	221 65	246 05	270 45
2: DNA pre-wash buffer	100	334 0	534 0	734 0	934 0	113 40	133 40	153 40	173 40	193 40	213 40	233 40
3: g-DNA wash buffer	100	534 0	934 0	133 40	173 40	213 40	253 40	293 40	333 40	373 40	413 40	453 40
4: Tris-HCl	10	702	902	110 2	130 2	150 2	170 2	190 2	210 2	230 2	250 2	270 2

3 Run the epMotion protocol 88x Quick-DNA Magbead Plus Low Volume 40_6.

3.1 Run the protocol with level sensing enabled.