VERSION 2

NOV 12, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.88ihzue

Protocol Citation: Monica Berjon-Otero, Sarah Duponchel, Matthias Fischer 2019. Lysozyme-based removal of bacteria from cultures of the marine heterotrophic flagellate *Cafeteria roenbergensis*. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.88ihzue>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Nov 12, 2019

Last Modified: Nov 12, 2019

PROTOCOL integer ID:
 29674

Lysozyme-based removal of bacteria from cultures of the marine heterotrophic flagellate *Cafeteria roenbergensis* V.2

Monica Berjon-Otero¹, Sarah Duponchel¹, Matthias Fischer¹

¹Max Planck Institute for Medical Research

Protist Research to Optimize Tools in Genetics (PROT-G)

Monica Berjon-Otero

ABSTRACT

This protocol allows to remove bacteria from cultures of the marine heterotrophic nanoflagellate *Cafeteria roenbergensis* without affecting flagellate viability. Bacterial removal is essential to reduce contamination and background signal in subsequent microscopy analysis, protein extraction, DNA extraction or other applications. With some changes, this protocol could be adapted to different types of protists.

MATERIALS

MATERIALS

- ☒ PBS **Contributed by users**
- ☒ MgCl₂ **Contributed by users**
- ☒ NaCl **Contributed by users**
- ☒ BD Bacto™ Yeast Extract **BD Biosciences Catalog #212750**
- ☒ OptiPrep™ 250 mL **Stemcell Technologies Catalog #7820**
- ☒ 2 mL Eppendorf **Contributed by users**
- ☒ Falcon tube (50 mL) **Contributed by users**
- ☒ DNase I (100 mg) **Roche Catalog #1284932**
- ☒ Thermomixer **Contributed by users**
- ☒ Lysozyme **Thermo Fisher Scientific Catalog #89833**
- ☒ High speed centrifuge **Contributed by users**
- ☒ Phase-contrast microscope **Contributed by users**
- ☒ F/2 artificial seawater medium **Contributed by users**
- ☒ Lugol's acid iodine solution **Contributed by users**
- ☒ Haemocytometer **Contributed by users**
- ☒ SW28 tubes 25x89 mm **Beckman Coulter Catalog #344058**

Preparation of *Cafeteria roenbergensis* culture

- 1 Determine the cell density of *Cafeteria roenbergensis*: stain 10 μL of a *Cafeteria* culture with 1 μL of Lugol's acid iodine solution and count the cells on a haemocytometer (Neubauer Chamber)
- 2 Dilute the *Cafeteria* culture to 5×10^5 cells/mL with f/2 medium + 0.03% (w/v) yeast extract. Incubate overnight at 22-25 °C.
- 3 Determine the flagellate concentration and use approx. 8×10^7 *Cafeteria* cells/sample
- 4 Centrifuge flagellates at 4,500 g, 5 min, room temperature in 50 mL Falcon tubes. Flagellates will be in the pellet.
- 5 Resuspend the cell pellets in 1.8 mL of 1X PBS and transfer them into 2 mL microfuge tubes.

Lysozyme and DNaseI treatment

- 6 Add 40 μL of 50 mg/mL **freshly prepared** lysozyme in 1X PBS (1 mg/mL final concentration). Incubate for 1 hour at room temperature.

Note

Optionally, you can also add 5 mM EDTA here. EDTA increases the activity of lysozyme against Gram negative bacteria, but also negatively affects *Cafeteria* mobility.

- 7 Add 200 μL of 10 mg/mL **freshly prepared** DNaseI in 1X PBS (1 mg/mL final concentration, 2

U/μL) and 11 μL of 1M MgCl₂ (5 mM final concentration).
Incubate for 30 min at 34°C in a thermomixer with shaking.

- 8 Centrifuge at 5,000 g, 6 min, 20°C

Optiprep purification

- 9 Resuspend the pellets in 100 μL of 1X gradient buffer (0.5 M NaCl in 1X PBS). Pipette up and down several times to break up aggregates.
- 10 Up to 1×10^9 *Cafeteria* cells can be loaded per SW28 centrifuge tube. Add 1X gradient buffer up to 0.5 - 1 mL final resuspended cell volume.
- 11 Load the sample on top of a two-step Optiprep cushion, 20% (w/v) Optiprep at the bottom of the tube and 10% (w/v) Optiprep on top of the 20% layer (both Optiprep solutions are prepared in 1X gradient buffer).
- 12 Centrifuge the tubes in an ultracentrifuge, Beckman SW28 rotor, at 20,000 g, 20 min, 20°C, **with slow braking**.
- 13 Recover *Cafeteria* cells from the gradient, at the interphase between the 10% and 20% Optiprep layers.

Recovery of *Cafeteria* culture

- 14 Dilute the recovered *Cafeteria* cells at least twofold with F/2 medium (no yeast extract)

15 Dilute the culture 25fold or 50fold with F/2 and stain 10 µL of this dilution with 1 µL of Lugol's acid iodine solution

Load the sample on a haemocytometer to quantify *Cafeteria* cells and remaining bacteria.

16 Dilute *Cafeteria* cells to the desired density. If no or little dilution is required, you can centrifuge again for 6 min at 5,000 g, 20°C and resuspend the cells in F/2 to remove all the remaining Optiprep.