

Jul 24, 2019 Version 1

## Make NGM medium V.1

DOI

[dx.doi.org/10.17504/protocols.io.5svg6e6](https://dx.doi.org/10.17504/protocols.io.5svg6e6)

Cancer Research UK / Wellcome Trust Gurdon Institute media kitchen<sup>1</sup>

<sup>1</sup>Cancer Research UK / Wellcome Trust Gurdon Institute



**Cristian Riccio**

University of Cambridge

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.5svg6e6](https://dx.doi.org/10.17504/protocols.io.5svg6e6)

**Protocol Citation:** Cancer Research UK / Wellcome Trust Gurdon Institute media kitchen 2019. Make NGM medium. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.5svg6e6>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**We use this protocol and it's working**


**Created:** July 24, 2019


**Last Modified:** July 24, 2019


**Protocol Integer ID:** 26165

## Materials


### MATERIALS

 double distilled water (ddH<sub>2</sub>O)

 Sodium chloride meets analytical specification of Ph.Eur Fisher Chemical **Fisher Scientific Catalog #S/3160/65**

 Agar **Formedium Catalog #AGA02**

 Bacto™ Peptone **Thermo Fisher Scientific Catalog #211677**

- 1  NGM\_agar\_5L w&p.xls
- 2 Measure approx 4.8L double distilled H2O in 5L bell jar with a magnetic flea.
- 3 Add bacto peptone and NaCl
- 4 Stir until all solutes are dissolved.
- 5 Dispense approx 972ml NGM media per 1L bottle OR 486ml NGM into a 500ml bottle.
- 6 Dispense 17g of agar and a magnetic flea to each 1L bottle OR 8.5g of agar and a magnetic flea to each 500ml bottle.
- 7 Label, date and autoclave.

8

NaCl		15g
bacto peptone	12.5g	
Double distilled H2O	4.86L	
Agar	per 1L bottle	17g