OCT 18, 2019

Making LB agar plates

Priota Islam¹

¹Imperial College London

Behavioural Genomics

Priota Islam

Imperial College London

ABSTRACT

Luria broth (LB) is a nutrient-rich media commonly used to culture bacteria in the lab. The addition of agar to LB results in the formation of a gel that bacteria can grow on, as they are unable to digest the agar but can gather nutrition from the LB within. When required the addition of an antibiotic to this gel allows for the selection of only those bacteria with resistance to that antibiotic - usually conferred by a plasmid carrying the antibiotic resistance gene.

See the link for more details: <https://www.addgene.org/protocols/pouring-lb-agar-plates/>

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.8eqhtdw

Protocol Citation: Priota Islam 2019. Making LB agar plates. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.8eqhtdw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 18, 2019

Last Modified: Oct 18, 2019

PROTOCOL integer ID:
28848

- 1 Purchase LB Agar from the media kitchen and autoclave it to melt

Content of LB Agar:

37 g pre-mixed powder consisting of:
- 5 g peptone

- 10 g peptone from casein
- 10 g sodium chloride
- 10 g agar-agar
- 1 L Sterile H₂O

2 If the bacteria have a specific antibiotic resistance then add 1000x of that antibiotic to the melted LB agar before pouring, e.g. if we use ampicillin we add 100mg/ml to the LB in a way that the final concentration becomes 100µg/ml. So, for 200ml LB agar we add 200ul of 100mg/ul of stock ampicillin

Antibiotic	Recommended Stock Concentration	Recommended Working Concentration
Ampicillin	100 mg/mL	100 µg/mL
Bleocin	5 mg/mL	5 µg/mL
Carbenicillin	100 mg/mL	100 µg/mL
Chloramphenicol	25 mg/mL (dissolve in EtOH)	25 µg/mL
Coumestrolin	25 mg/mL (dissolve in DMSO)	25 µg/mL
Gentamicin	10 mg/mL	10 µg/mL
Kanamycin	50 mg/mL	50 µg/mL
Spectinomycin	50 mg/mL	50 µg/mL
Tetracycline	10 mg/mL	10 µg/mL

Antibiotic Concentrations

- 3** Maintaining sterile techniques, pour the molten agar on large petri dishes (90mm) roughly about 30-45ml (don't need to be exact on the volume) and leave to dry overnight
- 4** Store the plates at 4C to be used later, ideally it should be used within 2 weeks but can be used for longer if not contaminated or dehydrated but never more than 3months