NOV 23, 2018

Making OP50 solution from Frozen Stock

Priota Islam¹¹Imperial College London

Behavioural Genomics

Priota Islam

Imperial College London

OPEN ACCESS**DOI:**dx.doi.org/10.17504/protocols.io.vube6sn

Protocol Citation: Priota Islam 2018. Making OP50 solution from Frozen Stock.

protocols.io<https://dx.doi.org/10.17504/protocols.io.vube6sn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Nov 23, 2018**Last Modified:** Nov 23, 2018**PROTOCOL integer ID:**

18019

¹ Day: 1

- Get LB Agar from the media kitchen and autoclave it for 2hrs
- Post autoclave, pour the agar on large petri dishes (60mm) and leave to dry overnight to be transferred to the cold room the next day

2 **Day: 2**

- Get the frozen tube out of the freezer (Freezer 12, SD Box)
- Take a sterile pipette tip and get some frozen sample
- Streak the pipette on the LB agar plate
- Take another new pipette and streak across the old streak and then at a separate spot streak again
- Repeat this step for one more time
- Keep the plate at 37°C incubator overnight for the bacteria to grow
- Transfer the plate to the fridge the next day (This plate can be used for about 1 month to inoculate bacterial culture)

3 **Day:3**

- Purchase LB Broth from the Media kitchen (Don't use less than 200ml)
- Get flat bottomed conical flasks from the glassware kitchen (For 200ml LB Broth take 400/500ml flask to allow enough headspace for the bacteria)
- From the culture plate, select and circle a single colony
- Using aseptic technique, carefully scoop the single colony and mix it with the LB Broth that is already poured into the conical flask
- Label the flask and put it on the 37°C shaker for overnight incubation

4 **Day:4**

- Following overnight incubation, take the flask out of the incubator and measure the OD using a spectrophotometer
- Record the average of the OD600, use LB Broth as Blank
- Aliquot the solution into labelled 15ml falcon tubes and keep them in the 4°C Fridge