VERSION 2
JUN 05, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.3qzgm6

Protocol Citation: Kenneth Schackart 2019. Mammalian Cell Culture: Refreshing Media. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.3qzgm6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jun 04, 2019

Last Modified: Jun 05, 2019

PROTOCOL integer ID:
24057

Mammalian Cell Culture: Refreshing Media V.2

Version 1 is forked from [Mammalian Cell Culture: Refreshing Media](#)

Kenneth
Schackart¹

¹University of Arizona

Yoon Lab

Kenneth Schackart
University of Arizona

ABSTRACT

This protocol explains how to refresh the media of cultured mammalian cells grown in a tissue culture flask.

GUIDELINES

- Gloves must be worn at all times.
- Perform all tasks within biosafety cabinet.
- Anything entering biosafety cabinet must be generously sprayed with 70% ethanol (even you).
- When finished, wipe biosafety cabinet with 70% ethanol, and UV for at least 15 minutes.

MATERIALS

- Gloves
- Cultured T-75 flask [or T-25 flask]
- Serological pipet with tips
- Waste beaker
- Dulbecco's Phosphate Buffered Saline (DPBS)
- Cell culture media (e.g. DMEM:F12, EMEM, etc.)

BEFORE START INSTRUCTIONS

- Warm cell culture media and DPBS in 37 °C water bath.
- Wash waste beaker with soap and warm water, then dry with paper towel.
- Expose serological pipet tips and waste beaker to UV for at least 00:15:00 .

Assessing Culture Health

- 1 Assess cell health under light microscope before beginning.

Refreshing media

- 2 Using a serological pipet, remove old media from flask and transfer to waste beaker.

Note

To avoid contamination, avoid touching the pipette tip to anything. Holding the flask such that the lid is pointed up will make it easier to remove all the liquid.

- 3 Wash cells by pipetting 4 mL warmed DPBS into the flask. [1 mL for T-25 flask]
- 4 Remove DPBS and dispose into waste beaker.
- 5 Repeat the above two steps for a total of 2 washes.
- 6 Pipette 8 mL - 12 mL warmed cell culture media into flask. [4 mL for T-25 flask].

Incubate

- 7 Spray flask generously with 70% ethanol solution before placing in CO₂ incubator.