JUN 05, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.x97fr9n

Protocol Citation: Kenneth Schackart 2019. Mammalian Cell Staining. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.x97fr9n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Feb 18, 2019

Last Modified: Jun 05, 2019

PROTOCOL integer ID:
 20511

Mammalian Cell Staining

Forked from [Mammalian Cell Staining](#)

Kenneth
 Schackart¹

¹University of Arizona

Yoon Lab

Kenneth Schackart
 University of Arizona

ABSTRACT

This protocol details how to stain mammalian cells cultured on a 96-well plate. Actin filaments, focal adhesion sites (as indicated by the presence of vinculin), and nuclei will be stained.

GUIDELINES

- Gloves must be worn at all times.
- Perform all tasks within biosafety cabinet.
- Anything entering biosafety cabinet must be generously sprayed with 70% ethanol (even you).
- When finished, wipe biosafety cabinet with 70% ethanol, and UV for at least 15 minutes.

MATERIALS

- 4% Paraformaldehyde solution
- 0.1% Triton X-100 solution in PBS
- Blocking buffer (PBS + 1% bovine serum albumin)
- Anti-vinculin solution (1:500 in blocking buffer)
- TRITC and FITC-conjugated secondary antibody solution (1:1:248, TRITC:FITC:blocking buffer) referred to as FITC:TRITC, prepared in opaque centrifuge tube
- DAPI solution (1:249 DAPI:blockingbuffer) prepared in opaque centrifuge tube
- Phosphate buffered saline (PBS)
- Washing buffer (PBS + 0.05% Tween-20)

BEFORE START INSTRUCTIONS

- Warm DPBS in 37 °C water bath.
- Wash waste beaker with soap and warm water, then dry with paper towel.
- Expose waste beaker to UV for at least 00:15:00 .

Fix the cells

- 1 Remove cell culture media.

Note

You may wash twice with DPBS at this point.

- 2 Add 100 µL of 4 % volume paraformaldehyde solution.
- 3 Incubate for 00:15:00 .

Perforate cell membrane

- 4 Remove paraformaldehyde solution.
- 5 Wash twice with 100 µL washing buffer.
- 6 Add 100 µL of 0.1 % volume Triton X-100.

7 Incubate for 00:05:00 .

Block unspecific binding

8 Remove Triton X-100.

9 Wash twice with washing buffer.

10 Add 100 μ L blocking buffer.

11 Incubate for 00:30:00 .

Bind anti-vinculin to vinculin

12 Remove blocking buffer.

13 Wash twice with washing buffer.

14 Add 100 μ L of Anti-Vinculin and blocking buffer mixture.

15 Incubate for 01:00:00 .

Note

Increasing this duration can greatly increase FITC-conjugated secondary antibody binding. It is acceptable to place covered plate in refrigerator overnight. While waiting on this incubation, it is a good time to prepare the combined dye solution.

Stain actin filaments and focal adhesion sites

16 Remove anti-vinculin blocking buffer mixture.

17 Wash twice.

18 Add 100 μ L FITC:TRITC solution

19 Incubate for 00:30:00 .

Stain nucleus

20 Remove stains.

- 21 Wash twice with washing buffer.
- 22 Add 100 μ L of DAPI solution
- 23 Incubate for 00:05:00
- 24 Remove DAPI.
- 25 Add 100 μ L PBS to prevent dessication of cells during imaging/storage.