JUN 19, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.qf5dtq6

External link:
<https://www.janelia.org/project-team/flylight/protocols>

Protocol Citation: Geoffrey W. Meissner, Jonathan B. Grimm, Rebecca M. Johnston, Ben Sutcliffe, Julian Ng, Gregory S.X.E. Jefferis, Sebastian Cachero, Luke D. Lavis, Oz Malkesman 2018. MCFO IHC of Drosophila CNS. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.qf5dtq6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: May 25, 2018

Last Modified: Jun 19, 2018

MCFO IHC of Drosophila CNS

Forked from [MCFO Hybrid Chemical Tag & IHC for Adult CNS](#)

Geoffrey W. Meissner¹, Jonathan B. Grimm¹, Rebecca M. Johnston¹, Ben Sutcliffe², Julian Ng², Gregory S.X.E. Jefferis², Sebastian Cachero², Luke D. Lavis¹, Oz Malkesman¹

¹Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, Virginia, USA;

²Division of Neurobiology, Medical Research Council Laboratory of Molecular Biology, Cambridge CB2 0QH, United Kingdom

Geoffrey Meissner

ABSTRACT

The use of genetically encoded 'self-labeling tags' with chemical fluorophore ligands enables rapid labeling of specific cells in neural tissue. To improve the chemical tagging of neurons, we synthesized and evaluated new fluorophore ligands based on Cy, Janelia Fluor, Alexa Fluor, and ATTO dyes and tested these with recently improved *Drosophila melanogaster* transgenes. We found that tissue clearing and mounting in DPX substantially improves signal quality when combined with specific non-cyanine fluorophores. We compared and combined this labeling technique with standard immunohistochemistry in the Drosophila brain.

GUIDELINES

- All tissues and solutions are at room temperature (RT), unless noted. Always protect tissue from light exposure.
- For details on dissection and fixation see FlyLight Protocol - Adult Dissection and 2% Fixation.
- For mounting and embedding instructions refer to FlyLight Protocol – DPX Mounting.
- Protocols at <https://www.janelia.org/project-team/flylight/protocols>
- **Reporter Genotype:** pBPhsFlp2::PEST in attP3; brp-SNAP / CyO; pJFRC201-10XUAS-FRT>STOP>FRT-myr::smGFP-HA in VK0005, pJFRC240-10XUAS-FRT>STOP>FRT-myr::smGFP-V5-THS-10XUAS-FRT>STOP>FRT-myr::smGFP-FLAG in su(Hw)attP1
- For details on reporter constructs see Nern, et al., 2015. <http://www.pnas.org/content/112/22/E2967.long> doi: 10.1073/pnas.1506763112

Reagents and Supplies

- AF594 Donkey α - Jackson Immuno Research. # 711-585-152
- ATTO 647N Goat α -Rat IgG (H&L) Antibody. Rockland. # 612-156-120
- DL550 Mouse α -V5 Tag. AbD Serotec. # MCA1360D550GA
- DPX Mountant for Microscopy. Electron Microscopy Sciences. # 13512, 500 mL
- Ethanol, ACS reagent, >99.5% (200 proof). Sigma Aldrich. # 459844-1L
- GS – Goat Serum. Life Technologies. 16210-064, 100 mL
- Kodak Photo-Flo 200 Solution. Electron Microscopy Sciences. # 74257
- nc82 – Mouse α - Developmental Studies Hybridoma Bank. # nc82-s
- Cy2 Goat α - Jackson Immuno Research. # 115-225-166
- NMS – Normal Mouse Serum. Jackson Immuno Research. # 015-000-120
- PBS – Phosphate Buffered Saline, 1X. # 21-040
- PFA – Paraformaldehyde. 20% PFA. Electron Microscopy Sciences. # 15713-S
- Poly-L-Lysine. Sigma Aldrich. # P1524-25MG
- Protein LoBind Microcentrifuge Tubes - 2 mL. # 022431102
- S2 – Schneider's Insect Medium. Sigma Aldrich. # S01416
- Rabbit α -HA Tag. Cell Signal Technologies. # 3724S
- Rat α -FLAG Tag (DYKDDDDK Epitope Tag). Novus Biologicals. # NBP1-06712
- Triton X-100. Sigma Aldrich. # X100
- Xylenes. Fisher Scientific. # X5-500

- 1 **Dissect.** Dissect adult brains or CNS in cold Schneider's Insect Medium (S2).
- 2 **Fix.** Transfer tissue to 2 mL Protein LoBind tubes filled with 2% paraformaldehyde (PFA) in S2 at RT. Fix for 55 minutes at RT while nutating.
- 3 **Post-fix wash.** Remove the fix and add 1.75 mL phosphate buffered saline with 0.5% Triton X-100 (PBT) and wash for a total of 4 X 10-minutes washes while nutating. If needed, store tissue in 0.5% PBT at 4°C while nutating or lay tube flat and rotate.
- 4 **Block Goat Serum (GS).** Remove PBT and add 200 μ L 5% GS in PBT per tube. Incubate for 1.5 hours at RT on a rotator with tubes upright.

- 5 Primary antibodies.** Remove block and add primary antibodies diluted in 5% GS in PBT for a volume of 200 μ L per tube. Incubate for 4 hours at RT on a rotator with tubes upright. Then continue incubation at 4°C on a rotator with tubes upright for 2 overnights.
- Mouse nc82 (1:30 or 33.3 μ L/mL)
Rat α -FLAG Tag (1:200 or 5 μ L/mL)
Rabbit α -HA Tag (1:300 or 3.3 μ L/mL)
- 6 Post-primary washes.** Remove the primary antibody and do a brief rinse with 1.75 mL 0.5% PBT. Allow the tissue to settle to the bottom and then remove the rinse solution and add 1.75 mL 0.5% PBT. Wash for a total of 5 X 15-minute washes while nutating.
- 7 Secondary antibodies.** Remove PBT and add the secondary antibodies diluted in 5% GS in PBT for a volume of 200 μ L per tube. Incubate for 4 hours at RT on a rotator with tubes upright. Then continue incubation at 4°C on a rotator with tubes upright for 3-4 overnights.
- Cy2 Goat α -Mouse (1:600 or 1.67 μ L/mL)
ATTO647N Goat α -Rat (1:300 or 3.3 μ L/mL)
AF594 Donkey α -Rabbit (1:500 or 2 μ L/mL)
- 8 Post-secondary washes.** Remove the secondary antibody and do a brief rinse with 1.75 mL 0.5% PBT. Allow the tissue to settle to the bottom and then remove the rinse solution and add 1.75 mL 0.5% PBT. Wash for a total of 5 X 15-minute washes while nutating. If needed, store tissue in 0.5% PBT at 4°C while nutating or lay tube flat and rotate.
- 9 Block Normal Mouse Serum (NMS).** Remove PBT and add 200 μ L of 5% NMS in PBT per tube. Incubate for 1.5 hours at RT on a rotator with tubes upright
- 10 Direct Label α -V5 antibody.** Remove NMS block and add DL550 Mouse α -V5 in 5% NMS in PBT. Incubate for 4 hours at RT on a rotator with tubes upright. Then continue incubation at 4°C on a rotator with tubes upright for 1 overnight.
- DL550 Mouse α -V5 (1:500 or 2 μ L/mL)
- 11 Post- α -V5 washes.** Remove the α -V5 antibody and do a brief rinse with 1.75 mL 0.5% PBT. Allow the tissue to settle to the bottom and then remove the rinse solution and add 1.75 mL 0.5% PBT. Wash for a total of 5 X 15-minute washes while nutating.
- 12 Pre-embedding fixation.** Remove PBT and add 1.75 mL 4% PFA in PBS at RT. Fix for 4 hours at RT while nutating.

- 13 Post-4% PFA washes.** Remove the 4% PFA and do a brief rinse with 1.75 mL 0.5% PBT. Allow the tissue to settle to the bottom and then remove the rinse solution and add 1.75 mL 0.5% PBT. Wash for a total of 4 X 15-minute washes while nutating. If needed, store tissue in 0.5% PBT at 4°C while nutating or lay tube flat and rotate.
- 14 Mount.** Mount the tissue on a poly-L-lysine (PLL) coated cover glass. • For making PLL see FlyLight Recipe – Poly-L-Lysine.
- 15 Dehydrate.** Move the cover glass through a series of 7 cover glass staining jars filled with increasing concentrations of ethanol (30%, 50%, 75%, 95%, 100%, 100%, 100%). Soak the cover glass for 10 minutes in each jar.
- 16 Xylene clearing.** (In the hood). Move the cover glass through a series of 3 jars filled with xylene. Soak the cover glass for 5 minutes in each jar.
- 17 DPX embedding.** Add 7 drops of dibutyl phthalate in xylene (DPX) on top of the tissue mounted on the cover glass. Place the cover glass (DPX down) on a prepared slide with spacers. Use the edge of a glass slide to gently press down on the center of the cover glass to seat the cover glass onto the slide. Let the slide dry in the hood for 2 days before viewing.