Molecular Crowding
SCR-seq

VERSION 2

MAY 22, 2018

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.p9kdr4w

External link:

<https://www.nature.com/articles/s41467-018-05347-6>

Protocol Citation: Johannes JWB Bagnoli, Christoph Ziegenhain, Aleksandar Janjic, Lucas Esteban Wange, Beate Vieth, Swati Parekh, Johanna Geuder, Ines Hellmann, Wolfgang Enard 2018. mcSCRB-seq protocol.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.p9kdr4w>

MANUSCRIPT CITATION:

Johannes W. Bagnoli, et al (2018) Sensitive and powerful single-cell RNA sequencing using mcSCRB-seq. *Nature Communications* 9:2937. doi: [10.1038/s41467-018-05347-6](https://doi.org/10.1038/s41467-018-05347-6)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

mcSCRB-seq protocol V.2

Nature Communications

Johannes JWB Bagnoli¹, Christoph Ziegenhain¹, Aleksandar Janjic¹, Lucas Esteban Wange¹, Beate Vieth¹, Swati Parekh¹, Johanna Geuder¹, Ines Hellmann¹, Wolfgang Enard¹

¹Ludwig-Maximilians-Universität München

Human Cell Atlas Method Development Community

Aleksandar Janjic

Ludwig-Maximilians-Universität München

ABSTRACT

Single-cell RNA sequencing (scRNA-seq) has emerged as a central genome-wide method to characterize cellular identities and processes. Consequently, improving its sensitivity, flexibility and cost-efficiency can advance many research questions. Among the flexible plate-based methods, “Single-Cell RNA-Barcoding and Sequencing” (SCR-seq) is one of the most sensitive and efficient ones. Here, we systematically evaluated experimental conditions of this protocol and find that adding polyethylene glycol considerably increases sensitivity by enhancing cDNA synthesis. Furthermore, using Terra polymerase increases efficiency due to a more even cDNA amplification that requires less sequencing of libraries. We combined these and other improvements to a new scRNA-seq library protocol we call “molecular crowding SCR-seq” (mcSCRB-seq), which we show to be the most sensitive and one of the most efficient and flexible scRNA-seq methods to date.

ATTACHMENTS

[mcSCRBseq_oligodT.txt](#)

Protocol status: Working
We use this protocol and it's working

Created: May 21, 2018

Last Modified: Jul 26, 2018

PROTOCOL integer ID:
12300

Keywords: molecular crowding, scRNA-seq, SCRB-seq

GUIDELINES

- For troubleshooting help, feel free to join our [mcSCRB-seq Slack channel](#), leave your question in the comments section, or message us directly.
- The complete list of reagents and plastic ware with order numbers can be found in the 'Materials' section.
- Make sure all steps involving single-cell lysate and RNA before reverse transcription are carried out swiftly.
- Size selection of libraries is optional, but has in our experience improved reliability in cluster densities when sequencing.
- All primer sequences are listed below:

Oligo	Vendor	Purification	Concentration	Sequence
barcoded oligo-dT (E3V6NEXT)	IDT	TruGrade	2 µM	Biotin-ACACTCTTTCCCTACACGACGCTCTTCCGATCT[B C6][UMI10][T30]VN
TSO unblocked (E5V6NEXT)	IDT	HPLC	100 µM	ACACTCTTTCCCTACACGACGCrGrGrG
PreAmp (SINGV6)	IDT	Desalted	10 µM	Biotin-ACACTCTTTCCCTACACGACGC
3' enrichment primer (P5NEXTPT 5)	IDT	HPLC	5 µM	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCG*A*T*C*T
i7 Index Primer (N7XX)	IDT	TruGrade	5 µM	CAAGCAGAAGACGGCATACGAGAT[i7]GTCTCGTGGGCTCGG

Find the cell barcode sequences in the attached text file (Abstract tab).

MATERIALS

MATERIALS

- ⊗ EDTA 0.5M **Sigma Aldrich Catalog #E7889**
- ⊗ Ethanol, absolute **Carl Roth Catalog #9065.4**
- ⊗ Exonuclease I (20 U/µl) **Thermo Fisher Scientific Catalog #EN0582**
- ⊗ Exonuclease I Reaction Buffer (10x) **Thermo Fisher Scientific Catalog #EN0582**

✕	IGEPAL CA-630 Sigma Aldrich Catalog #I8896
✕	Maxima H- Reverse Transcriptase (200 U/μl) Thermo Fisher Scientific Catalog #EP0753
✕	Maxima RT Buffer (5x) Thermo Fisher Scientific Catalog #EP0753
✕	Polyethylene glycol 8000 Sigma Aldrich Catalog #89510
✕	PBS 7.4 Gibco - Thermo Fischer Catalog #10010-23
✕	Phusion HF Buffer New England Biolabs Catalog #B0518
✕	Proteinase K Takara Catalog #9034
✕	Sera-Mag Speed Beads Thermo Fisher Scientific Catalog #65152105050250
✕	Terra PCR Direct Polymerase Mix Takara Catalog #639271
✕	UltraPure DNase/RNase-Free Distilled Water Invitrogen - Thermo Fisher Catalog #10977-049
✕	Sodium Azide 99.5% Sigma Aldrich Catalog #S2002-100G
✕	Sodium Chloride 5M Sigma Aldrich Catalog #S5150-1L
✕	Trizma hydrochloride solution 1M pH 8.0 Sigma Aldrich Catalog #T2694
✕	Bioanalyzer High Sensitivity DNA Analysis Kits Agilent Technologies Catalog #5067-4626
✕	MinElute Gel Extraction Kit Qiagen Catalog #28606
✕	Nextera XT DNA Library Preparation Kit illumina Catalog #FC-131-1096
✕	Quant-iT PicoGreen dsDNA Assay Kit Invitrogen - Thermo Fisher Catalog #P7589
✕	dNTPs (25 mM each) Thermo Fisher Scientific Catalog #R0182
✕	Aluminium seals for cold storage Contributed by users Catalog #391-1275
✕	Adhesive PCR film seals Thermo Fisher Scientific Catalog #AB0558
✕	twin.tec 96-well DNA LoBind Plates Eppendorf Catalog #0030129504
✕	twin.tec 384-well DNA LoBind Plates Eppendorf Catalog #0030129547
✕	0.5 ml PCRclean tube DNA LoBind Eppendorf Catalog #0030108035
✕	1.5 ml PCRclean tube DNA LoBind Eppendorf Catalog #0030108051
✕	5.0 ml PCRclean tube DNA LoBind Eppendorf Catalog #0030108310
✕	15 ml PCRclean tube DNA LoBind Eppendorf Catalog #0030122208
✕	E-Gel EX Agarose Gels, 2% Invitrogen - Thermo Fisher Catalog #G402002

BEFORE START INSTRUCTIONS

Wipe bench surfaces with RNase Away and keep working environment clean.

Preparation of lysis plates

- 1 Prepare **Lysis Buffer** according to the number of plates to be filled.

Reagent	96-well plate	384-well plate
NEB HF Phusion buffer (5x)	1.1 µL	4.4 µL
Proteinase K (20 mg/mL)	27.5 µL	110 µL
UltraPure Water	411.4 µL	1645.6 µL
Total	440 µL	1760 µL

- 2 Prepare 96/384 well plate(s) containing 4 µL **Lysis Buffer** per well.

Add 1 µL barcoded oligo-dT primer [2 µM] (E3V6NEXT adapter) to each well (12-/64-channel pipette).

Note

Lysis plates with barcode primers can be prepared ahead of time and stored at -20 °C

🧪 4 µL Lysis Buffer

🧪 1 µL barcoded oligo-dT primer [2 µM]

Sample Collection

- 3 Sort 1 cell to each well of a 96/384 well plate containing 5 µL **Lysis Buffer** and **barcoded oligo-dT primer**.
- 4 Immediately seal the plate with an aluminium cold storage seal.
- 5 In a cooled centrifuge, spin down the plate for 30 sec @ max. speed and place immediately on dry ice.

🌡️ -80 °C Store plates containing single-cell lysates in a -80 °C freezer for up to 6 months.

Proteinase K Digest

- 6 Thaw plates briefly (up to 1 min) at room temperature
- 7 Spin down (30 sec @ 1000 rcf) in a centrifuge pre-cooled to 4 °C.
🌡️ 4 °C
- 8 Replace aluminum foil seal with PCR plate seal to avoid excessive stickiness of the glue.
- 9 In a thermocycler with heated lid, incubate as follows:
🌡️ 50 °C 10 min (Proteinase K digest)
🌡️ 80 °C 10 min (Heat inactivation)
🌡️ 8 °C ∞ (Store)

Note

During incubation, proceed with preparation of **Reverse Transcription Mix**.

Reverse Transcription

- 10 Prepare **Reverse Transcription Mix** as follows:

Reagent	96-well plate	384-well plate
UltraPure Water	88 µL	352 µL
PEG 8000 (50 % solution)	165 µL	660 µL
Maxima RT Buffer (5x)	220 µL	880 µL
dNTPs (25 mM each)	44 µL	176 µL

TSO E5V6NEXT unblocked (100 μ M)	22 μ L	88 μ L
Maxima H Minus RT (200 U/ μ l)	11 μ L	44 μ L
Total	550 μL	2200 μL

Note

Caution: Reverse Transcription Mix with PEG needs to be mixed carefully!

4 °C Keep Reverse Transcription Mix on ice

11 Add 5 μ L Reverse Transcription Mix to each well.

5 μ L Reverse Transcription Mix

Note

If a robot (eg. Formulatrix Mantis) is used, make sure to calibrate correctly to the viscous solution.

12 Seal plate with a PCR seal, vortex briefly and spin down (30 sec @ 1000 rcf) in a centrifuge pre-cooled to 4 °C.

13 In a thermocycler with heated lid, incubate:

42 °C 90 min

8 °C ∞

cDNA Pooling & Purification

14 Prepare **Pooling Beads**:

Reagent	Amount
PEG 8000	15 g
NaCl, 5M	20 mL
Tris-HCL, 1M, pH 8.0	500 µL
EDTA, 0.5M	100 µL
IGEPAL, 10% solution	50 µL
Sodium Azide, 10% solution	250 µL
UltraPure Water	up to 50 mL
Total	50 mL

Add all ingredients into a 50 mL falcon tube, but do not add the total amount of water until after PEG is completely solubilized

Incubate at 40°C and vortex regularly until PEG is completely dissolved

Resuspend bead stock carefully (Sera-Mag Speed Beads)

Pipette 100 µL of bead suspension into a 96-well plate well

Place on magnet stand

Remove supernatant

Add 100 µL 10 mM Tris-HCl, pH 8.0, 1 mM EDTA (TE), and resuspend beads

Place on magnet stand

Remove supernatant

Repeat wash one more time

Add 90 µL 10 mM Tris-HCl, pH 8.0, 1 mM EDTA (TE), and resuspend beads

Add to PEG solution above and mix well.

Note

Beads can be prepared ahead of time and stored at 4 °C or room temperature.

Note

Caution: Beads tend to pellet and need to be carefully resuspended before use.

15 For 96-well plates: Pool all wells of one plate into a 2 mL tube and add 960 µL (ratio 1/1) **30% PEG Pooling Beads**

For 384-well plates: Pool all wells of one plate into a 15 mL falcon tube and add 3840 µL (ratio 1/1) **30% PEG Pooling Beads**

🧪 960 µL Pooling Beads (96-well plates)

🧪 3840 µL Pooling Beads (384-well plates)

16

00:05:00 binding of the cDNA onto the beads

20 °C Room temperature

- 17 Place on magnet stand until clear
For 384-well plates: Remove supernatant, leaving about 1 mL in the tube. Resuspend the beads in the leftover supernatant and transfer to a 1.5 mL tube for easier handling.

- 18 Discard supernatant

- 19 Wash twice with 1 mL 80% ethanol (while on magnet) and discard supernatant

1 mL 80% ethanol (freshly prepared)

- 20 00:05:00 air dry beads

- 21 Elute cDNA in 17 µL UltraPure Water & transfer to new tube

17 µL UltraPure Water

Exonuclease I Treatment

- 22 To the 17 µl cDNA, add:

2 µL Exonuclease I Buffer (10x)

1 µL Exonuclease I (20 U/µl)

- 23 In a thermocycler with heated lid, incubate:

37 °C 20 min (ExoI digest)

🔥 80 °C 10 min (Heat inactivation)

🧊 8 °C ∞ (Store)

Full length cDNA amplification

24 Prepare **PreAmplification Mix** as follows:

Reagent	1x
Terra direct Buffer (2x)	25 µL
SINGV6 Primer (10 µM)	1 µL
Terra polymerase (1.25 U/µL)	1 µL
UltraPure Water	3 µL
Total	30 µL

25 Add 30 µL of **PreAmplification Mix** directly to the Exonuclease I digested sample.

🧊 30 µL PreAmplification Mix

26 In a thermocycler with heated lid, incubate as follows:

Step	Temperature	Time	Cycles
Initial Denaturation	98 °C	3 min	1x
Denaturation	98 °C	15 sec	13-21x
Annealing	65 °C	30 sec	
Elongation	68 °C	4 min	
Final Elongation	72 °C	10 min	1x
Store	8 °C	∞	

Note

Cycle number highly depends on the input amount and should be optimized depending on the specific celltype used in the experiment. For ES cells, 13-15 cycles are sufficient.

cDNA purification & quantification

27 Prepare **Clean-up Beads**:

Reagent	Amount
PEG 8000	11 g
NaCl, 5M	10 mL
Tris-HCL, 1M, pH 8.0	500 µL
EDTA, 0.5M	100 µL
IGEPAL, 10% solution	50 µL
Sodium Azide, 10% solution	250 µL
UltraPure Water	up to 49 mL
Total	49 mL

Add all ingredients into a 50 mL falcon tube, but do not add the total amount of water until after PEG is completely solubilized

Incubate at 40°C and vortex regularly until PEG is completely dissolved

Resuspend bead stock carefully (Sera-Mag Speed Beads)

Pipette 1000 µL of bead suspension into a 1.5 mL tube

Place on magnet stand

Remove supernatant

Add 1000 µL 10 mM Tris-HCl, pH 8.0, 1 mM EDTA (TE), and resuspend beads

Place on magnet stand

Remove supernatant

Repeat wash one more time

Add 900 µL 10 mM Tris-HCl, pH 8.0, 1 mM EDTA (TE), and resuspend beads

Add to PEG solution above and mix well.

Note

Beads can be prepared ahead of time and stored at 4 °C or room temperature.

28 Mix PreAmplification PCR with 40 µL **Clean-up Beads** (1/0.8 ratio)

40 µL Clean-up Beads

29 00:05:00 binding of the cDNA onto the beads

- 30** Place on magnet until clear and discard supernatant
- 31** Wash twice with 150 μ L 80% ethanol (while on magnet) and discard supernatant
150 μ L 80% ethanol (freshly prepared)
- 32** 00:05:00 air dry beads
- 33** Elute cDNA in 15 μ L UltraPure Water & transfer to new tube
15 μ L UltraPure Water
- 34** Quantify the cDNA using the Quant-iT PicoGreen dsDNA assay kit or equivalent Qubit following the manufacturer's protocol. Use 1 μ L of clean cDNA for quantification.

Expected result

cDNA concentration should be > 1 ng/ μ L, depending on cell type and cycle number

- 35** Optional: Quality check the cDNA using the Agilent 2100 Bioanalyzer with High Sensitivity DNA Analysis Kits.

Expected result

Tagmentation, Library PCR & Indexing

- 36** Prepare **Tagmentation Mix** and dispense 19 μL to a new 96-well plate.

Reagent	1x
Tagment DNA Buffer (2x)	10 μL
Amplicon Tagment Mix (Tn5)	5 μL
UltraPure Water	4 μL
Total	19 μL

🧴 19 μL Tagmentation Mix

- 37** Dilute cDNA to 0.8 ng/ μL and add 1 μL to each reaction.

🧴 1 μL cDNA (0.8 ng/ μL)

38 In a thermocycler with heated lid, incubate as follows:

55 °C Tagmentation

00:10:00 Tagmentation

39 To stop the reaction, add 5 µL NT buffer to each reaction and mix by pipetting up and down.

5 µL NT Buffer

00:05:00 Incubation at room temperature

40 Prepare **3' Enrichment PCR Mix** as follows and add 24.5 µL to each tagmentation reaction.

Reagent	1x
NPM PCR Mix	15 µL
P5NEXTPT5 (5 µM)	0.5 µL
UltraPure Water	9 µL
Total	24.5 µL

24.5 µL 3' Enrichment PCR Mix

41 Add 0.5 µL of i7 index primer (5 µM)

0.5 µL i7 index primer (5 µM)

42 In a thermocycler with heated lid, incubate as follows:

Step	Temperature	Time	Cycles
Gap-fill	72 °C	3 min	1x
Initial Denaturation	95 °C	30 sec	
Denaturation	95 °C	10 sec	13x
Annealing	55 °C	30 sec	
Elongation	72 °C	1 min	
Final Elongation	72 °C	5 min	1x
Store	8 °C	∞	

43 Mix Index PCR with 50 µL **Clean-up Beads** (1/1 ratio)

🧪 50 µL Clean-up Beads

44 ⌚ 00:05:00 binding of DNA onto the beads

45 Place on magnet until clear and discard supernatant

46 Wash twice with 150 µL 80% ethanol (while on magnet) and discard supernatant

🧪 150 µL 80% ethanol (freshly prepared)

47 ⌚ 00:05:00 air dry beads

48 Elute cDNA in 20 µL UltraPure Water & transfer to new tube

🧪 20 µL UltraPure Water

Size selection

49 Load complete library onto an 2% Agarose E-Gel EX and run for 10 minutes.

⌚ 00:10:00

50 As soon as the Gel run has finished open the Gel framing using the Gel opening tool

51 Excise the Library from 300bp to 900bp using a clean scalpel

52 Gel purify the slice using the Qiagen MinElute Kit following manufacturer's guidelines:

- Add 450 µL Buffer QG
- Dissolve the gel slice in QG for 10 min @ 42 °C
- Add 150 µL Isopropanol to the sample and mix by inverting
- Transfer sample to spin column and centrifuge at 16 000 x g for 1 min
- Discard flow through and add 500 µL Buffer QG
- Centrifuge at 16 000 x g for 1 min and discard flow through
- Add 700 µL Buffer PE
- Centrifuge at 16 000 x g for 1 min and discard flow through
- Centrifuge again at 16 000 x g for 1 min to remove residual ethanol
- Transfer column to a new 1.5 mL microcentrifuge tube
- Add 20 µL H₂O to column and incubate for 1 min
- Centrifuge at 16 000 x g for 1 min to elute and discard the spin column

Library Quantification

53 Quantify and quality control the library using the Agilent 2100 Bioanalyzer with High Sensitivity DNA Analysis Kits.

Load both the library and a 1/10 dilution on two different lanes of the chip.

Note

If cDNA was not quality controlled previously, it is strongly suggested to run cDNA on the same chip.

Expected result

Successful libraries will typically exceed 3-5 ng/ μ l concentration.

- 54 If pooling several libraries, combine equal molar amounts.

Sequencing

- 55 Sequence your library on any compatible Illumina sequencer.
Dilute libraries to recommended molarity according to Illumina's recommendations (eg. 2 nM).
Select the following paired-end read-length settings:

	Read	Cycles	Content
	Read 1	16	Cell barcode & UMI
	Index 1	8	i7 Index
	Index 2	0	
	Read 2	50	cDNA fragment

Primary data processing using zUMIs

- 56** Download and install zUMIs including all dependencies.

Software

zUMIs	NAME
Linux	OS
https://github.com/sdparekh/zUMIs	SOURCE LINK

- 57** Copy the sequencing data from the sequencer and run bcl2fastq without demultiplexing.

Command

```
bcl2fastq --use-bases-mask Y16,I8,Y50 --create-fastq-for-index-reads
```

- 58** Run zUMIs with the following parameters. Replace Read names and paths to reference genome and annotation with actual files of your instance.

Command

```
zUMIs-master.sh -f lane1.R1.fastq.gz -c 1-6 -m 7-16 -T lane1.I1.fastq.gz -U 1-8 -r  
lane1.R2.fastq.gz -l 50 -n mcSCRBseqrun -p 16 -a /path_to/gene_annotation.gtf -g  
/path_to/reference_genome_index
```