VERSION 3

DEC 11, 2019

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.v63e9gn

External link:

<https://doi.org/10.7717/peerj.8157>

Protocol Citation: Kiichi

Hirota, Yoshiyuki Matsuo
2019. Measurement of insulin
concentration. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.v63e9gn>

MANUSCRIPT CITATION:

License: This is an open
access protocol distributed
under the terms of
the [Creative Commons
Attribution License](#), which
permits unrestricted use,
distribution, and reproduction
in any medium, provided the
original author and source are
credited

Protocol status: Working

Created: Dec 06, 2018

Last Modified: Dec 11, 2019

Measurement of insulin concentration V.3

PeerJ

Kiichi Hirota¹, Yoshiyuki Matsuo¹

¹Kansai Medical University

Yoshiyuki Matsuo

BEFORE START INSTRUCTIONS

Insulin secreted into the culture medium from MIN6 and INS-1 cells was measured using the Mouse and Rat Insulin H-type™ enzyme-linked immunosorbent assay kit (Shibayagi Co. Ltd., Shibukawa, Japan), according to the manufacturer's protocol (Suzuki K. et al. 2015, PeerJ 3:e1498. doi: 10.7717/peerj.1498).

- 1** MIN6 cells were cultured for 30 min in Krebs-Ringer bicarbonate HEPES (KRBH) buffer (140 mM NaCl, 3.6 mM KCl, 0.5 mM NaH₂PO₄, 0.5 mM MgSO₄, 1.5 mM CaCl₂, 2 mM NaHCO₃, 10 mM HEPES, and 0.1% bovine serum albumin) containing 2 mM glucose.
- 2** The buffer was then changed to another KRBH buffer solution containing the relevant experimental concentration of glucose, and the cells were cultured for 1 h.
- 3** The KRBH buffer was collected and its insulin concentration was measured.
- 4** The insulin concentration of the buffer was normalized to the total cell protein level.

The results were normalized to the concentration of control samples of each independent experiment and the normalized values were shown as insulin secretion ratio.