JAN 03, 2018

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.j9dcr26](https://doi.org/10.17504/protocols.io.j9dcr26)

External link:
<https://doi.org/10.1371/journal.pone.0190595>

Protocol Citation: Bianca Hemmeryckx, Dries Bauters, H. Roger Lijnen 2018. Metabolic parameters determination in acid citrated plasma. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.j9dcr26>

MANUSCRIPT CITATION: Bauters D, Bedossa P, Lijnen HR, Hemmeryckx B (2018) Functional role of ADAMTS5 in adiposity and metabolic health. PLoS ONE 13(1): e0190595. doi: [10.1371/journal.pone.0190595](https://doi.org/10.1371/journal.pone.0190595)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Metabolic parameters determination in acid citrated plasma

PLOS One

Dries
 Bianca Hemmeryckx¹, Bauters¹,
¹KU Leuven
 H. Roger Lijnen¹

Bianca Hemmeryckx

ABSTRACT

Diet-induced obesity by administration of a western-type diet high in fat and sugar to mice may result in an elevation of plasma total and LDL-cholesterol levels, triglycerides and the levels of the liver enzymes alkaline phosphatases and aspartate (AST) and alanine aminotransferases (ALT).

GUIDELINES

- The acid citrate solution is best kept at +4°C and made fresh every month.
- Concentrations of plasma metabolic parameters need to be multiplied by the dilution factor 3.

MATERIALS

STEP MATERIALS

- Trisodium citrate dihydrate Merck MilliporeSigma (Sigma-Aldrich) Catalog #S1804
- citric acid monohydrate Merck MilliporeSigma (Sigma-Aldrich) Catalog #33114
- Trisodium citrate dihydrate Merck MilliporeSigma (Sigma-Aldrich) Catalog #S1804
- citric acid monohydrate Merck MilliporeSigma (Sigma-Aldrich) Catalog #33114

Created: Oct 11, 2017

Last Modified: Feb 21, 2018

PROTOCOL integer ID:
8197

Keywords: Diet-induced
obesity, liver and metabolism

- 1 Collect 300µl blood from the retro-orbital sinus using a capillary tube containing acid citrate and collect the blood in an eppendorf tube containing 30µl acid citrate (1/10 dilution). Do not include the first two drops of blood.

Acid citrate (50 ml): dissolve 1.25g trisodiumcitrate dihydrate (85 mM) and 0.82g citric acid monohydrate (78 mM) in 20 ml milliQ water and add MilliQ water until volume is 50 ml. pH should be 4.5.

Trisodium citrate dihydrate Merck MilliporeSigma (Sigma-Aldrich) Catalog
#S1804

citric acid monohydrate Merck MilliporeSigma (Sigma-Aldrich) Catalog
#33114

- 2 Plasma is obtained by centrifuging the blood in an eppendorf centrifuge for 10 min at 10,000 rpm. Collect supernatant.
- 3 Mix 50µl of acid citrated plasma with 100µl metabolic buffer consisting of 0.1 M Tris pH 7.5 and 0.15 M NaCl in a 1 ml sample cup provided by the routine lab of UZ Leuven, Leuven, Belgium.
- 4 Transfer samples to the routine lab of UZ Leuven, Leuven, Belgium for the measurement of plasma levels of total cholesterol, LDL- and HDL-cholesterol, triglycerides, alkaline phosphatases and aspartate and alanine aminotransferases.