AUG 05, 2020

Metabolite Extraction

avinash.kale¹¹domnic colvin

UM-DAE Centre for Excellence in Basic Sciences

avinash.kale

OPEN ACCESS**DOI:**dx.doi.org/10.17504/protocols.io.bjdtki6n

Document Citation: avinash.kale 2020. Metabolite Extraction. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bjdtki6n>

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Aug 05, 2020**Last Modified:** Aug 05, 2020**DOCUMENT integer ID:**
40083**Metabolite Extraction**

The bacterial cultures were grown overnight in 200 ml LB media to sufficiently synthesize their metabolites. The bacterial cultures were centrifuged and washed to remove traces of remaining media. Cell pellets were then sonicated at amplitude 30%, pulse 10 seconds, gap 10 seconds, for a total time of 2 minutes: 30 seconds or 15 Cycles in 300 µL of 1x PBS. The samples were centrifuged at 12000 rpm for 20 minutes and the supernatant was separated. The metabolites were precipitated using acetonitrile in twice the volume of the supernatant by incubating at -80 °C overnight. The clear supernatant was centrifuged, transferred into new tubes, followed by air drying(27). These dried samples were further given for LCMS.