

Jan 11, 2020 Version 1

Methods and protocols from Chaibarat et al. (2019) for systematic reintroduction of bateng (*Bos javanicus*) V.1

DOI

dx.doi.org/10.17504/protocols.io.ba8wihxe

Rattanawat Chaibarat¹, Nampung Youngpoy², Praeploy Kongsurakan², Seree Nakbun³

¹Wildlife and Plants Research Center, Faculty of Environment and Resource Studies, Mahidol University;

²Wildlife and Plants Research Centre, Faculty of Environment and Resource Studies, Mahidol University;

³Khao Nampu Nature and Wildlife Education Center, Department of National Parks, Wildlife and Plant Conservation, Kanchanaburi, 71250, Thailand

WILDREC

Tech. support email: rattanawat.cha@mahidol.ac.th

[Click here to message tech. support](#)



Rattanawat Chaibarat

Wildlife and Plants Research Center, Faculty of Environment ...

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.ba8wihxe

External link: <http://DOI: 10.1071/WR18184>

Protocol Citation: Rattanawat Chaibarat, Nampung Youngpoy, Praeploy Kongsurakan, Seree Nakbun 2020. Methods and protocols from Chaibarat et al. (2019) for systematic reintroduction of bateng (*Bos javanicus*). **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.ba8wihxe>

Manuscript citation:

Chaibarat, R., Youngpoy, N., Kongsurakan, P. & Nakboon, S. (2019). Habitat preferences of reintroduced banteng (*Bos javanicus* d'Alton 1823) into the Salakphra Wildlife Sanctuary, Thailand. Wildl. Res. 46(7), 573-586. <http://DOI: 10.1071/WR18184>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working



Created: January 11, 2020

Last Modified: January 11, 2020

Protocol Integer ID: 31734

Keywords: Banteng, reintroduction, soft release, training protocols

Abstract

Banteng (*Bos javanicus*) underwent general medical checkups and received minimal human contact in 302 ha enclosure. They were trained in transportation boxes (1 m × 2.5 m × 1.8 m, width × long × high) over 6 months. They were kept in groups in soft release cage for 4 months. Natural food plants, fresh water and artificial salt licks were provided. Immobilizations were controlled with anesthetic drugs by veterinarians and fitted with radio collars. The radio signals were monitored every week through ground tracking, using homing in and triangulation techniques via VHF signals. Camera traps were installed, memory cards and batteries were changed every month in each location.

Attachments



[Protocol_bateng.docx](#)

113KB

