MojoSort™ Human NK Cell Isolation Kit Protocol

Sam Li¹

¹BioLegend

Sam Li
BioLegend

ABSTRACT

BioLegend MojoSort™ nanobeads work in commonly used separation columns, based on our internal research as well as validation by external testing by academic labs. This simple protocol consists of following the MojoSort™ protocol to label the cells with **pre-diluted** MojoSort™ reagents and using the columns as indicated by the manufacturer.

Note: Due to the properties of our beads, it may be possible to use far fewer beads and less antibody cocktail than with other commercial suppliers. We recommend a titration to find the best dilution factor. However, as a general rule, dilutions ranging from 1:2 to 1:10 for the antibody cocktail can be used. Dilutions ranging from 1:5 to 1:20 for the Streptavidin Nanobeads can be used. Please contact BioLegend Technical Service (tech@biolegend.com) if further assistance is needed.

GUIDELINES

MojoSort™ magnetic particles can be used with other commercially available magnetic separators, both free standing magnets and column-based systems. Because MojoSort™ protocols are optimized for the MojoSort™ separator, the protocols may need to be adjusted for other systems. Please contact BioLegend Technical Service (tech@biolegend.com) for more information and guidance. We do not recommend using MojoSort™ particles for BD's IMag™ or Life Technologies' DynaMag™.

MATERIALS

MATERIALS

MojoSort™ Buffer BioLegend Catalog
#480017

Additional reagents:

- commercially available cell separation columns
- 5 mL polypropylene tubes
- 70 µm cell strainer

This protocol works with the following MojoSort™ Kits (cat#):
[480053](#), [480054](#)

DOI:
[dx.doi.org/10.17504/protocols.io.7bihike](https://doi.org/10.17504/protocols.io.7bihike)

External link:
<https://www.biolegend.com/protocols/mojosort-human-nk-cell-isolation-kit-column-protocol/4622/>

Protocol Citation: Sam Li .
MojoSort™ Human NK Cell Isolation Kit Protocol.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.7bihike>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Sep 13, 2019

Last Modified: Sep 13, 2019

PROTOCOL integer ID:
27722

Keywords: MojoSort™ , nanobeads, column, cell separation

1 Prepare cells from your tissue of interest or blood without lysing erythrocytes.

- 2 In the final wash of your sample preparation, resuspend the cells in MojoSort™ Buffer by adding up to 4 mL in a 5 mL (12 x 75 mm) polypropylene tube.
Note: Keep MojoSort™ Buffer on ice throughout the procedure.
- 3 Filter the cells with a 70 µm cell strainer, centrifuge at 300xg for 5 minutes, and resuspend in an appropriate volume of MojoSort™ Buffer. Count and adjust the cell concentration to 1×10^8 cells/mL. 5m
- 4 Aliquot 100 µL (10^7 cells) into a new tube. Add **10 µL of the pre-diluted Biotin-Antibody Cocktail**. Mix well and **incubate on ice for 15 minutes**. Scale up the volume if separating more cells. For example, add 100 µL of pre-diluted antibody cocktail for separating 1×10^8 cells in 1 ml of MojoSort™ Buffer. When working with less than 10^7 cells, use indicated volumes for 10^7 cells. 15m
- 5 Wash the cells by adding MojoSort™ Buffer up to 4 mL. Centrifuge the cells at 300xg for 5 minutes. 5m
- 6 Discard the supernatant and resuspend cells in 100 µL of MojoSort™ Buffer.
- 7 Vortex the Streptavidin conjugated Nanobeads (to resuspend) at max speed, 5 touches, and prepare the dilutions to test. Add **10 µL of pre-diluted Streptavidin Nanobeads**. Mix well and **incubate on ice for 15 minutes**. Scale up the volume accordingly if separating more cells. For example, add 100 µL of pre-diluted Nanobeads for separating 1×10^8 cells in 1 ml of MojoSort™ Buffer. When working with less than 10^7 cells, use indicated volumes for 10^7 cells. 15m
- 8 Wash the cells by adding MojoSort™ Buffer up to 4 mL. Centrifuge the cells at 300xg for 5 minutes. 5m
- 9 Discard the supernatant.
- 10 Resuspend the cells in appropriate amount of MojoSort™ Buffer and proceed to separation. At least 500µL is needed for column separation.

Note: There are several types of commercially available columns, depending on your application. Choose the one that fits best your experiment:

	Max. number of labeled cells	Max. number of total cells	Cell suspension volume	Column rinse volume	Cell wash volume	Elution volume
Small Capacity	1×10^7	2×10^8	500 μ L for up to 10^8 cells	1ml	1 ml	1 ml
Medium Capacity	1×10^8	2×10^9	500 μ L for up to 10^9 cells	3ml	3 ml	5 ml
Large Capacity	1×10^9	2×10^{10}	500 μ L for up to 10^{10} cells	20-50ml	30 ml	20 ml

Example of magnetic separation with medium capacity column

- 11 Place the column in a magnetic separator that fits the column.
- 12 Rinse the column with 3 mL of cell separation buffer.
- 13 Add the labeled cell suspension in at least 500 μ L of buffer to the column through a 30 μ m filter and collect the fraction containing the unlabeled cells. These are the cells of interest; do not discard.
- 14 Wash the cells in the column **1 time** with 3 mL of buffer and collect the fraction containing the unlabeled cells. Combine with the collected fraction from step 3.
- 15 If desired, the labeled cells can be collected by taking away the column from the magnet and place it on a tube. Then add 5 mL of buffer and flush out the magnetically labeled fraction with a plunger or supplied device. The labeled cells may be useful as staining controls, to monitor purity/yield, or other purposes.