MojoSort™ Mouse anti-APC Nanobeads Protocol

Sam Li[1]

[1]BioLegend

BioLegend
Tech. support email: **tech@biolegend.com**

Sam Li
BioLegend

DOI:
dx.doi.org/10.17504/protocols.io.7xhhpj6

External link:
https://www.biolegend.com/protocols/mojosort-mouse-anti-apc-nanobeads-protocol/4756/

Protocol Citation: Sam Li . MojoSort™ Mouse anti-APC Nanobeads Protocol. **protocols.io** https://dx.doi.org/10.17504/protocols.io.7xhhpj6

License: This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Oct 03, 2019

Last Modified: Oct 03, 2019

PROTOCOL integer ID: 28361

ABSTRACT

Product description and procedure summary: Target cells are positively selected or depleted by incubating the sample with an anti-human APC conjugated antibody, followed by incubation with magnetic anti-APC Nanobeads. The magnetically labeled fraction is retained by the use of a magnetic separator. These are the APC+ cells, do not discard them if those are the cells of interest. Some of the downstream applications include functional assays, gene expression, phenotypic characterization, etc.

Note: This procedure is optimized for the isolation of 10^7 to 2 x 10^8 cells per tube. If working with fewer than 10^7 cells, keep volumes as indicated for 10^7 cells. For best results, optimize the conditions to your specific cell number and tissue. Prepare fresh MojoSort™ Buffer solution by diluting the 5X concentrate with sterile distilled water.*Scale up volumes if using 14mL tubes and Magnet, and place the tube in the magnet for 10 minutes.*

GUIDELINES

MojoSort™ magnetic particles can be used with other commercially available magnetic separators, both free standing magnets and column-based systems. Because MojoSort™ protocols are optimized for the MojoSort™ separator, the protocols may need to be adjusted for other systems. Please contact BioLegend Technical Service (tech@biolegend.com) for more information and guidance. We do not recommend using MojoSort™ particles for BD's IMag™ or Life Technologies' DynaMag™.

Application notes: To use this product in magnetic separation columns, a titration of the Nanobeads should be performed. Optimal concentration for magnetic separation columns is lot-specific. Please contact BioLegend Technical Service (tech@biolegend.com) for further assistance on how to use MojoSort™ Nanobeads in magnetic separation columns.

Keywords: mojosort, anti-APC, nanobeads, isolation, selection, magnet

MATERIALS

MATERIALS

MojoSort™ Magnet **BioLegend Catalog #**480019

MojoSort™ Buffer **BioLegend Catalog #**480017

MojoSort™ Mouse anti-APC Nanobeads **BioLegend Catalog #**480071, 480072

- Adjustable pipettes
- 70µm filters (one per sample)
- 5mL (12 x 75mm) or 14mL (17 x 100mm) polypropylene tubes polypropylene tubes
- Reagents for sample preparation
- Reagents and instruments (Flow cytometer) to determine yield and purity

1 Prepare cells from your tissue of interest or blood without lysing erythrocytes.

2 In the final wash of your sample preparation, resuspend the cells in MojoSort™ Buffer by adding up to 4 mL in a 5 mL (12 x 75 mm) polypropylene tube.
Note: Keep MojoSort™ Buffer on ice throughout the procedure.

3 Filter the cells with a 70µm cell strainer, centrifuge at 300xg for 5 minutes, and resuspend in an appropriate volume of MojoSort™ Buffer. Count and adjust the cell concentration to 1 x 10^8 cells/mL.

4 Aliquot 100µL of cell suspension (10^7 cells) into a new tube. Check the recommended usage for flow cytometric staining of the APC-conjugated antibody indicated in the antibody technical datasheet. Calculate the volume to stain 10^7 cells (or desired amount of cells). **Add the appropriate volume of APC-conjugated antibody** to the cell suspension, mix well and **incubate on ice for 15 minutes**.
Optional: Take an aliquot before adding the cocktail to monitor purity and yield.

5 Wash the cells by adding MojoSort™ Buffer up to 4mL. Centrifuge the cells at 300xg for 5 minutes.

6 Discard supernatant and resuspend in 100µL of MojoSort™ Buffer.

7 Resuspend the beads by vortexing, maximum speed, 5 touches. Add **10µL of APC Nanobeads.** Mix well and **incubate on ice for 15 minutes**. Scale up the volume accordingly if separating more cells. For example, add 100 µL of Nanobeads for separating 1 x 10^8 cells in 1 ml of MojoSort™ Buffer. When working with less than 10^7 cells, use indicated volumes for 10^7 cells.

8 Wash the cells by adding MojoSort™ Buffer up to 4mL. Centrifuge the cells at 300xg for 5 minutes.

9 Discard the supernatant.

10 Add 2.5mL of MojoSort™ Buffer.
Note: If you observe aggregates, filter the suspension. To maximize yield, you can disrupt the aggregates by pipetting the solution up and down.

11 Place the tube in the magnet for 5 minutes.
Optional: Take a small aliquot before placing the tube in the magnet to monitor purity and yield. Keep unused cells to be used as control or other applications if needed.

12 Pour out the unlabeled fraction. If these are your cells of interest, **DO NOT DISCARD**. Resuspend the labeled cells in 2.5mL MojoSort™ Buffer.

13 Repeat steps 10-12 on the labeled fraction twice more for a total of **3 separations.** Pool the unlabeled fractions and keep the labeled cells. The fraction that is not of interest may be useful as staining controls, to monitor purity/yield, or other purposes.
Optional: Take a small aliquot to monitor purity and yield.


Antibody
15 min
on ice
Nanobeads
Wash
15 min
on ice
Buffer
Wash
5 min in
magnet
MojoSort™
BioLegend®
Collect labeled cells,
resuspend in buffer
Repeat twice with
labeled cell fraction for a
total of 3 separations
Cells labeled with
Nanobeads
Pour off unlabeled cells