

## MojoSort™ Selection Kits Protocol - 5

DOI

[dx.doi.org/10.17504/protocols.io.7z9hp96](https://dx.doi.org/10.17504/protocols.io.7z9hp96)



Sam Li<sup>1</sup>

<sup>1</sup>BioLegend

BioLegend

Tech. support email: [tech@biolegend.com](mailto:tech@biolegend.com)



Sam Li

BioLegend



**DOI:** [dx.doi.org/10.17504/protocols.io.7z9hp96](https://dx.doi.org/10.17504/protocols.io.7z9hp96)

**External link:** <https://www.biolegend.com/protocols/mojosort-selection-kits-protocol-5/4244/>

**Protocol Citation:** Sam Li . MojoSort™ Selection Kits Protocol - 5. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.7z9hp96>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Created:** October 08, 2019

**Last Modified:** October 08, 2019

**Protocol Integer ID:** 28449

**Keywords:** mojosort, selection, isolation, magnet

## Abstract

### Product description and procedure summary:

Target cells are either selected or depleted by incubating the sample with the biotin antibody cocktail followed by incubation with magnetic Streptavidin Nanobeads (Cat. No. 480015/480016). The magnetically labeled fraction is retained by the use of a magnetic separator. The untouched cells are collected. If these are the cells of interest; do not discard the liquid. Some of the downstream applications include functional assays, gene expression, phenotypic characterization, etc.

### Sample Preparation:

Enzymatic digestion of mouse brain followed by myelin removal is recommended to achieve the highest purity and yield. There are several protocols published that can be applied. As a general guideline, Trypsin digestion followed by a 70/37/30% percoll gradient will increase final purity and yield. *Scale up volumes if using 14 ml tubes and Magnet, and place the tube in the magnet for 10 minutes.*

## Guidelines


**Important Note:** MojoSort™ magnetic particles can be used with other commercially available magnetic separators, both free standing magnets and column-based systems. Because MojoSort™ protocols are optimized for the MojoSort™ separator, the protocols may need to be adjusted for other systems. Please contact BioLegend Technical Service (tech@biolegend.com) for more information and guidance. We do not recommend using MojoSort™ particles for BD's IMag™ or Life Technologies' DynaMag™.

**Application notes:** To use this product in magnetic separation columns, a titration of the Nanobeads should be performed. Optimal concentration for magnetic separation columns is lot-specific. Please contact BioLegend Technical Service (tech@biolegend.com) for further assistance on how to use MojoSort™ Nanobeads in magnetic separation columns.


## Materials

### MATERIALS

 MojoSort™ Buffer **BioLegend Catalog #480017**

 MojoSort™ Magnet **BioLegend Catalog #480019**

 MojoSort™ Mouse CX3CR1 Selection Kit **BioLegend Catalog #480055, 480056**

 MojoSort™ Mouse P2RY12 Selection Kit **BioLegend Catalog #480113, 480114**

- Adjustable pipettes
- 70µm filters (one per sample)
- 5mL (12 × 75mm) or 14mL (17 × 100mm) polypropylene tubes
- Reagents for sample preparation
- Reagents and instruments (Flow cytometer) to determine yield and purity



- 1 Prepare cells from your tissue of interest or blood without lysing erythrocytes.
- 2 In the final wash of your sample preparation, resuspend the cells in MojoSort™ Buffer by adding up to 4 mL in a 5 mL (12 × 75 mm) polypropylene tube.  
**Note:** Keep MojoSort™ Buffer on ice throughout the procedure.
- 3 Filter the cells with a 70µm cell strainer, centrifuge at 300xg for 5 minutes, and resuspend in an appropriate volume of MojoSort™ Buffer. Count and adjust the cell concentration to  $1 \times 10^7$  cells/mL.
- 4 Aliquot 100µL of cell suspension ( $10^6$  cells) into a new tube. Add **10µL of the Biotin-Antibody Cocktail**. Mix well and **incubate on ice for 15 minutes**. Scale up the volume accordingly if separating more cells. For example, add 100 µL of Antibody Cocktail for separating  $1 \times 10^7$  cells in 1 ml of MojoSort™ Buffer. When working with less than  $10^6$  cells, use indicated volumes for  $10^6$  cells.  
*Optional: Take an aliquot before adding the cocktail to monitor purity and yield.*
- 5 Wash the cells by adding MojoSort™ Buffer up to 4mL. Centrifuge the cells at 300xg for 5 minutes.
- 6 Discard supernatant and resuspend in 100µL of MojoSort™ Buffer.
- 7 Resuspend the beads by vortexing, maximum speed, 5 touches. Add **10µL of Streptavidin Nanobeads**. Mix well and **incubate on ice for 15 minutes**. Scale up the volume accordingly if separating more cells. For example, add 100 µL of Nanobeads for separating  $1 \times 10^7$  cells in 1 ml of MojoSort™ Buffer. When working with less than  $10^6$  cells, use indicated volumes for  $10^6$  cells.
- 8 Wash the cells by adding MojoSort™ Buffer up to 4mL. Centrifuge the cells at 300xg for 5 minutes.
- 9 Discard supernatant.
- 10 Add 2.5mL of MojoSort™ Buffer.  
**Note:** If you observe aggregates, filter the suspension. To maximize yield, you can disrupt the aggregates by pipetting the solution up and down.
- 11 Place the tube in the magnet for 5 minutes.  
*Optional: Take a small aliquot before placing the tube in the magnet to monitor purity and yield. Keep unused cells to be used as control or other applications if needed.*

- 12 Pour out the unlabeled fraction. If these are your cells of interest, **DO NOT DISCARD**. Resuspend the labeled cells in 2.5mL MojoSort™ Buffer.
- 13 Repeat steps 10-12 on the labeled fraction twice more for a total of **3 separations**. Pool the unlabeled fractions and keep the labeled cells. The fraction that is not of interest may be useful as staining controls, to monitor purity/yield, or other purposes.  
*Optional: Take a small aliquot to monitor purity and yield.*

