JUN 24, 2020

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.bgcajsse](https://doi.org/10.17504/protocols.io.bgcajsse)

External link:
<https://doi.org/10.1371/journal.pone.0234958>

Protocol Citation: Hossein Tarrahimofrad, Amir Meimandipour, Sareh Arjmand, Mohammadtaghi Beigi Nassiri, Ehsan Jahangirian, Hossein Tavana, Javad Zamani, Somayyeh Rahimnahal, Saeed Aminzadeh 2020. Molecular dynamics simulation (Protein-Ligand). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bgcajsse>

MANUSCRIPT CITATION: Tarrahimofrad H, Meimandipour A, Arjmand S, Nassiri MB, Jahangirian E, Tavana H, Zamani J, Rahimnahal S, Aminzadeh S (2020) Structural and biochemical characterization of a novel thermophilic Coh01147 protease. PLoS ONE 15(6): e0234958. doi: [10.1371/journal.pone.0234958](https://doi.org/10.1371/journal.pone.0234958)

Molecular dynamics simulation (Protein-Ligand)

PLOS One

Hossein Tarrahimofrad¹, Amir Meimandipour¹, Sareh Arjmand², Mohammadtaghi Beigi Nassiri³, Ehsan Jahangirian¹, Hossein Tavana⁴, Javad Zamani¹, Somayyeh Rahimnahal¹, Saeed Aminzadeh¹

¹Bioprocess engineering group, Institute of Industrial and Environmental Biotechnology, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran;

²Protein Research Center, Shahid Beheshti University, G. C., Tehran, Iran;

³Department of Animal Science and Food Technology, Agriculture Science and Natural Resources University Khuzestan, Ahwaz, Iran;

⁴Department of Biomedical Engineering, The University of Akron, Akron, OH 44325, USA

Hossein Tarrahimofrad

ABSTRACT

The prepared protein and protein-ligand complexes (protease-SDS and protease-tween complexes) were used in MD simulation, performed using GROMACS v 4.6.5 with CHARMM36 all-atom force field. The CGenFF server (<https://cgenff.umaryland.edu/>) provides topologies and parameters of ligands compatible with the CHARMM36 all-atoms force field. Protein and protein-ligand complexes were soaked in a cubic box of water molecules, and the charges on the protein were neutralized by the addition of Na⁺ and Cl⁻ ions. The energy of the system was minimised using the steepest descent algorithm to eliminate bad contact and clashes. The NVT and NPT ensembles were used during the equilibration to achieve the desired temperature (373.15 K for protease and 300 K for protease-ligand complex) and pressure (1 bar) for 100 ps and restraint forces of 1000 kJ/mol. Finally, 40 ns MD run were performed in triplicate after releasing all restraints. All bonds were constrained by the LINCS algorithm.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: May 13, 2020

Last Modified: Jun 24, 2020

PROTOCOL integer ID:
36962