JUL 12, 2019

Mouse Stellate Immunohistochemistry protocol

Pradeep Rajendran¹

¹[University of California, Los Angeles]

Scott John

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.2qggdtw

Protocol Citation: Pradeep Rajendran 2019. Mouse Stellate Immunohistochemistry protocol. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.2qggdtw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 08, 2019

Last Modified: Jul 12, 2019

PROTOCOL integer ID:
23016

1 See mouse stellate isolation protocol

- 2 Complete/intact stellate ganglia are fixed in 4% paraformaldehyde overnight at 4°C.
- 3 Fixed tissue is rinsed in phosphate buffered saline (PBS), and stored in PBS + 0.02% sodium azide.
- 4 The stellate ganglion whole mounts are permeabilized in 'block' solution. (10% normal donkey serum in 1.0% BSA + 0.4% Triton X-100 + PBS).
- 5 Ganglia are stained with antibodies directed to either tyrosine hydroxylase (TH) or protein gene product 9.5 (PGP 9.5). Secondary staining with streptavidin conjugated to ATTO-647N was used to visualize neurobiotin filling. Stained tissue was rinsed in PBS and mounted on glass slides.