

Oct 24, 2019

Mowiol mounting media

DOI

dx.doi.org/10.17504/protocols.io.8p6hvre

Michael Economo¹

¹Boston University



Michael Economo

Boston University

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.8p6hvre

External link: <http://cshprotocols.cshlp.org/content/2010/1/pdb.rec12110.full>

Protocol Citation: Michael Economo 2019. Mowiol mounting media. [protocols.io](#)

<https://dx.doi.org/10.17504/protocols.io.8p6hvre>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 24, 2019

Last Modified: October 24, 2019

Protocol Integer ID: 29150

Keywords: histology, mounting media, embedding media

Abstract

This makes a low cost mounting media that preserves many fluorescent dyes very well. It is a non-hardening media.

This protocol first published in Cold Spring Harbor Protocols.

Materials

MATERIALS

 Glycerol Merck MilliporeSigma (Sigma-Aldrich) Catalog #G5516

 0.2 M Tris Buffer pH 8.5 Merck MilliporeSigma (Sigma-Aldrich) Catalog #T1503-100G

 Mowiol® 4-88 Merck MilliporeSigma (Sigma-Aldrich) Catalog #81381-50G

- 1 1.Mix glycerol and Mowiol; dissolve with frequent agitation for 1 h at room temperature.
- 2 2.Add H₂O. Stir for 1 h at room temperature.
- 3 3.Add Tris-Cl. Incubate for 2 h at 50°C with occasional stirring (e.g., for 2 min every 20 min).
- 4 4.Centrifuge at 5000g for 15 min.
- 5 5.Store 3-mL aliquots of the supernatant for up to 12 mo at -20°C.
Mowiol often does not dissolve completely.