

Nov 15, 2022

MTT assay

DOI

dx.doi.org/10.17504/protocols.io.ewov1oq9ylr2/v1

PMAT0001¹

¹Lee Kong Chian School of Medicine

LKC Translational Neuro



PMAT PMAT0001

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.ewov1oq9ylr2/v1

Protocol Citation: PMAT0001 2022. MTT assay. protocols.io <https://dx.doi.org/10.17504/protocols.io.ewov1oq9ylr2/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: November 15, 2022

Last Modified: November 15, 2022

Protocol Integer ID: 72745

Abstract

The MTT assay is based on the conversion of MTT (3-[4,5-dimethylthiazol-2-yl]-2,5 diphenyl tetrazolium bromide) into purple formazan crystals by LIVING cells, which determines the mitochondrial activity of these cells. Viable cells contain NADPH-dependent oxidoreductase enzymes which reduce MTT to formazan.¹ The insoluble formazan crystals are dissolved using a solubilization solution (e.g. acidified isopropanol, DMSO) and the resultant coloured solution is quantified by measuring the absorbance at 500-600nm using a multi-plate spectrophotometer. The darker the solution, the greater the number of viable, metabolically active cells.² Since for most cell populations, the total mitochondrial activity would be proportional to the number of viable cells, this assay is widely used in characterizing the cytotoxic effects of drugs or nanoparticle formulations on immortalized cell line or primary cell cultures.

Day 0







- 1 Passage cells which are at 80-90% confluence. Perform the usual trypsinization procedure.
- 2 Seed 15,000 cells well (100uL) in complete DMEM (using 1X DMEM) in a 96 well plate. Make sure to fill the peripheral wells with 100uL of water. Only use the inner 60 wells for test wells. Seed cells in one plate solely for the test concentrations (10 concentrations, sextuplicates). Seed another plate just to keep nanoparticle-only, DMEM-only and negative controls.


Day 1

- 3 Remove expired media.
- 4 Without any PBS wash, replace the wells with 100uL of nanoparticle-containing media of different concentrations. Use complete medium for this step (more serum-free or reduced serum media)

Day 2

2h 16m 25s

- 5 Remove ALL media
- 6 Replace with 100uL of complete DMEM + MTT (MTT final concentration = 0.5mg/mL in DMEM)
- 7 After adding DMEM +MTT, shake the plates for  00:00:30 at 70rpm on an orbital shaker. 30s
- 8 Incubate wells with MTT for  02:00:00 at  37 °C 2h
- 9 Remove all media + MTT from wells
- 10 Add 100uL DMSO into all the wells and shake for  00:00:10 at 70rpm on an orbital shaker. 10s
- 11 Incubate for  00:15:00 at  37 °C 15m

- 12 Place the plates in a microplane reader. Shake the plate using the in-built shaker feature for at least  00:00:45 at 425 cpm (fast) using double orbital shaking method. Following that, take absorbance measurements of wells @ 570nm using a microplane reader

45s