



Jul 27, 2022

Multiplex PCR Assay for Clade-typing Salmonella Enteritidis

DOI

<https://dx.doi.org/10.17504/protocols.io.4r3l2ok1jv1y/v1>

Sarah Gallichan^{1,2,3}, Blanca M. Perez-Sepulveda⁴, Nicholas A. Feasey^{3,5}, Jay C. D. Hinton⁴, Anthony M. Smith^{1,2}

¹Centre for Enteric Diseases at the National Institute for Communicable Diseases (NICD), Johannesburg, South Africa;

²Department of Clinical Microbiology and Infectious Diseases, School of Pathology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa;

³Department of Clinical Sciences, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, UK;

⁴Clinical Infection, Microbiology & Immunology, Institute of Infection, Veterinary & Ecological Sciences (IVES), University of Liverpool, Liverpool, United Kingdom;

⁵Malawi Liverpool Wellcome Research Programme, Kamuzu University of Health Sciences, Blantyre, Malawi



Sarah Gallichan

Liverpool School of Tropical Medicine

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.4r3l2ok1jv1y/v1>

Protocol Citation: Sarah Gallichan, Blanca M. Perez-Sepulveda, Nicholas A. Feasey, Jay C. D. Hinton, Anthony M. Smith 2022. Multiplex PCR Assay for Clade-typing Salmonella Enteritidis. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.4r3l2ok1jv1y/v1>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 21, 2022

Last Modified: July 27, 2022

Protocol Integer ID: 67219

Keywords: Non-typhoidal Salmonella, Real-time PCR, Phylogeny, Molecular surveillance, recognition of different salmonella enteritidis clade, different salmonella enteritidis clade, comparison of salmonella enteritidi, salmonella enteritidi, salmonella enteritidis salmonella enteritidi, typing salmonella enteritidis salmonella enteritidi, salmonella, multiplex pcr assay for clade, global epidemic clade, epidemiology of these clade, whole genome sequencing, genome sequencing, associated gastroenteritis, whole genome sequencing capacity, multiplex pcr assay, gastroenteritis, african clade, genome sequence, identification of these clade, named global epidemic, genome, epidemiology, clade, distinct clade, sequencing, global epidemic, causing human disease

Funders Acknowledgements:

German Federal Ministry of Education and Research

Grant ID: BMBF grant number: 81203616

Wellcome Trust Senior Investigator award

Grant ID: grant number: 106914/Z/15/Z

Disclaimer

In respect of the phyletic structure of *Salmonella* Enteritidis, we have designed primers to distinguish three clades and an outlier cluster in a single reaction. These are henceforth denoted "Regional" and "Clade". The purpose of the Regional (African or Global classification) and Clade (Global Epidemic, Global Outlier, East or West African classification) assays is to further classify *Salmonella* Enteritidis isolates to better understand the transmission and epidemiology of each *Salmonella* Enteritidis clade. The Regional and Clade assays described here are limited to previously confirmed *Salmonella* Enteritidis isolates.

Abstract

Salmonella Enteritidis is one of the most commonly reported serovars of non-typhoidal *Salmonella* causing human disease and is responsible for both gastroenteritis and invasive non-typhoidal *Salmonella* (iNTS) disease worldwide. Whole-genome sequence (WGS) comparison of *Salmonella* Enteritidis isolates from across the world have identified three distinct clades, named Global Epidemic, Central/East African and West African, all of which have been implicated in epidemics: the Global Epidemic clade was linked to poultry-associated gastroenteritis, while the two African clades were related to iNTS disease. Despite the recognition of different *Salmonella* Enteritidis clades, the distribution and epidemiology of these clades across Africa is poorly understood because currently identification of these clades requires whole genome sequencing capacity. Here, we developed a sensitive, time- and cost-effective real-time PCR assay capable of differentiating between the *Salmonella* Enteritidis clades to facilitate surveillance and to inform public health response.

Attachments



[iivxbvu27.pdf](#)

81KB

Guidelines

In respect of the phyletic structure of *S. Enteritidis*, we have designed primers to distinguish three clades and an outlier cluster in a single reaction. These are henceforth denoted "Regional" and "Clade". The purpose of the Regional (African or Global classification) and Clade (Global Epidemic, Global Outlier, East or West African classification) assays is to further classify *S. Enteritidis* isolates to better understand the transmission and epidemiology of each *S. Enteritidis* clade. The Regional and Clade assays described here are limited to previously confirmed *S. Enteritidis* isolates.

References:

1. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, et al. Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics*. 2015 Nov 15;31(22):3691–3.
2. Rutledge RG, Co C. Mathematics of quantitative kinetic PCR and the application of standard curves.

Preparation of control panel isolates

- 1 Select twelve *Salmonella* Enteritidis isolates that represent the Global Epidemic, Outlier, East African and West African clades (three biological replicates per clade) predicted by the hierBAPS (hierarchical Bayesian Analysis of Population Structure) algorithm.

Citation

Feasey NA, Hadfield J, Keddy KH, Dallman TJ, Jacobs J, Deng X, Wigley P, Barquist L, Langridge GC, Feltwell T, Harris SR, Mather AE, Fookes M, Aslett M, Msefula C, Kariuki S, MacLennan CA, Onsare RS, Weill FX, Le Hello S, Smith AM, McClelland M, Desai P, Parry CM, Cheesbrough J, French N, Campos J, Chabalgoity JA, Betancor L, Hopkins KL, Nair S, Humphrey TJ, Lunguya O, Cogan TA, Tapia MD, Sow SO, Tennant SM, Bornstein K, Levine MM, Lacharme-Lora L, Everett DB, Kingsley RA, Parkhill J, Heyderman RS, Dougan G, Gordon MA, Thomson NR (2016) . Distinct *Salmonella* Enteritidis lineages associated with enterocolitis in high-income settings and invasive disease in low-income settings. Nature genetics.

<https://doi.org/10.1038/ng.3644>

LINK

- 2 Streak the twelve selected *Salmonella* Enteritidis isolates on 5% blood agar (Diagnostic Media Products, Johannesburg, South Africa) plates and incubate  Overnight in an IN 750 incubator (Mettler, Schwabach, Germany) at  37 °C .
- 3 Pick single colonies from the blood agar plates and resuspend in  400 µL of 10X TE buffer ( 800 mL distilled water,  2.92 g Tris,  15.76 g EDTA ( 8)) in 2 mL Safe-Lock tubes (Eppendorf, Hamburg, Germany).
- 4 Perform a genomic DNA extraction using the QIAamp DNA Mini Kit (QIAGEN, Hilden, Germany) according to the manufacturer's recommendations.



Designing primers and probes for the Regional- and Clade-typing assays



- 5 Analyse the annotated whole genome sequences for the twelve selected *Salmonella* Enteritidis isolates using ROARY v.3.11.2, to identify genes that can uniquely distinguish the geographical region (Global, which includes the Global and Outlier clade, and African, which includes the East and West African clades) and each clade

Citation

Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush D, Keane JA, Parkhill J (2015). Roary: rapid large-scale prokaryote pan genome analysis. Bioinformatics (Oxford, England).

<https://doi.org/10.1093/bioinformatics/btv421>

LINK

- 6 Use EnteroBase v1.1.3 (accessible online: https://enterobase.warwick.ac.uk/species/senterica/search_strains) Multi locus sequence query to confirm in silico that the selected genes can classify sequenced *Salmonella* Enteritidis isolates into clades based on the presence and absence of the selected genes.
- 7 Design primer and probe sets using the online PrimerQuest tool (Integrated DNA Technology; accessible online: <https://eu.idtdna.com/pages/tools/primerquest>) using sequences from the selected genes.
- 8 Prepare primer and probe sets according to the manufacturer's recommendations.

Real-time PCR set up

- 9 Dilute primer and probe sets to a concentration of [M] 20 millimolar (mM) using nuclease-free water (Ambion, ThermoFisher Scientific, California, USA).
- 10 Create two master mixes, one with the primer and probe sets for the Regional assay (to distinguish between the Global and African regions) and one with the primer and probe sets for the Clade assay (to distinguish between the Global Epidemic, Global Outlier, East and West African).
- 11 In a 96-well reaction plate (Applied Biosystems, ThermoFisher Scientific, California, USA), place  25 µL of TaqMan Gene Expression Master Mix (ThermoFisher

Scientific, California, USA),  17.8 μL of nuclease-free water (Ambion, ThermoFisher Scientific, California, USA),  3 μL of the relevant Master Mix (depending on whether the Regional/ Clade assay is preformed) and  1.2 μL of DNA template in each well.

- 12 Seal the reaction plate with MicroAmp® Optical Adhesive Film (Applied Biosystems, Life Technologies™, California, USA) and centrifuged at  15000 rpm for  00:01:00 using an Allegra™ X-22R Centriuge (Beckman Coulter™, California, USA) to ensure all reagents are concentrated at the bottom of the wells.

1m



- 13 Load the reaction plate into a 7500 Real Time PCR System (Applied Biosystems, Life Technologies™, California, USA) and set up using the following reaction conditions:
 50 °C for  00:02:00 , followed by  95 °C for  00:10:00 and 40 cycles of  95 °C for  00:00:15 ,  60 °C for  00:00:30 and  72 °C for  00:00:30 .

13m 15s



Assessing performance of the Regional- and Clade-typing assays

- 14 Preform 10-fold serial dilutions on the genomic DNA from two control panel Salmonella Enteritidis isolates that together contain all the target genes.
- 15 Quantify the DNA concentrations spectroscopically using a NanoDrop 1000 Spectrophotometer (ThermoFisher Scientific, California, USA).
- 16 Set up a real-time PCR assay as described above.
- 17 Determine the limit of detection by assessing the DNA concentration yielding the highest Ct (cycle threshold) value under 30 cycles.
- 18 Calculate the linear range for the Ct values of the Regional- and Clade-typing assays using the CORREL function in Microsoft Excel 2010.
- 19 Plot a calibration curve (depicting the change in cycle threshold value with the change in log DNA concentration) and calculate the amplification efficiency (PCR efficiency = $10^{-1/\text{slope} - 1}$)



Citation

Gevertz JL, Dunn SM, Roth CM (2005). Mathematical model of real-time PCR kinetics. Biotechnology and bioengineering.

Citations

Step 1

Feasey NA, Hadfield J, Keddy KH, Dallman TJ, Jacobs J, Deng X, Wigley P, Barquist L, Langridge GC, Feltwell T, Harris SR, Mather AE, Fookes M, Aslett M, Msefula C, Kariuki S, Maclennan CA, Onsare RS, Weill FX, Le Hello S, Smith AM, McClelland M, Desai P, Parry CM, Cheesbrough J, French N, Campos J, Chabalgoity JA, Betancor L, Hopkins KL, Nair S, Humphrey TJ, Lunguya O, Cogan TA, Tapia MD, Sow SO, Tennant SM, Bornstein K, Levine MM, Lacharme-Lora L, Everett DB, Kingsley RA, Parkhill J, Heyderman RS, Dougan G, Gordon MA, Thomson NR. Distinct Salmonella Enteritidis lineages associated with enterocolitis in high-income settings and invasive disease in low-income settings.

<https://doi.org/10.1038/ng.3644>

Step 19

Gevertz JL, Dunn SM, Roth CM. Mathematical model of real-time PCR kinetics.

Step 5

Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush D, Keane JA, Parkhill J. Roary: rapid large-scale prokaryote pan genome analysis.

<https://doi.org/10.1093/bioinformatics/btv421>