JUN 25, 2021

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.buaynsfw

Protocol Citation: Marcus A A Woodworth, Hao Yuan Kueh, Joshua C. Vaughan 2021. Multiplexed, single-cell profiling of histone modifications with SCEPTRE. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.buaynsfw>

MANUSCRIPT CITATION: Woodworth, M.A., Ng, K.K.H., Halpern, A.R., Pease, N.A., Nguyen, P.H.B., Kueh, H.Y. and Vaughan, J.C. (2021) Multiplexed single-cell profiling of chromatin states at genomic loci by expansion microscopy. *Nucleic Acids Research*, 10.1093/nar/gkab423.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol for several histone modifications and genomic loci within a few different cell lines.

Multiplexed, single-cell profiling of histone modifications with SCEPTRE

Marcus A A Woodworth¹, Hao Yuan Kueh²,
 Joshua C. Vaughan^{1,3}

¹Department of Chemistry, University of Washington, Seattle, Washington, 98195, USA;

²Department of Bioengineering, University of Washington, Seattle, Washington, 98195, USA;

³Department of Physiology and Biophysics, University of Washington, Seattle, Washington, 98195, USA

Human Cell Atlas Method Development Community

Marcus A A Woodworth

ABSTRACT

Genome architecture and activity are tightly regulated during an organism's development and function. Histone modifications are thought to contribute to this regulation by acting in combination to specify the activity states of individual genes. However, most methods used to profile these modifications either require a large number of cells or only target an individual histone modification at a time. This protocol uses the method Single Cell Evaluation of Post-TRANslational Epigenetic Encoding (SCEPTRE) to quantify immunolabeled H3K4me3 and H3K27me3 histone modifications at specific non-repetitive genomic loci in single cells using Expansion Microscopy. The protocol can be adapted to target alternative histone modifications or protein assemblies in other cell lines and for other genomic regions of interest.

Reference:

Woodworth, M.A., Ng, K.K.H., Halpern, A.R., Pease, N.A., Nguyen, P.H.B., Kueh, H.Y. and Vaughan, J.C. (2021) Multiplexed single-cell profiling of chromatin states at genomic loci by expansion microscopy. *Nucleic Acids Research*, 10.1093/nar/gkab423.

Created: Apr 17, 2021

Last Modified: Jun 25, 2021

PROTOCOL integer ID:
49208

Keywords: Histone modifications, Expansion Microscopy, Fluorescence in situ Hybridization, Single Cell Profiling

ATTACHMENTS

[GAPDH_oligo_spreadsheets.xlsx](#)

GUIDELINES

The Single Cell evaluation for Post-Translational Epigenetic Encoding (SCEPTRE) protocol is performed as follows:

1. Cells are grown and fixed on coverslips.
2. Histone modifications are immunolabeled.
3. Immunolabeled cells are gelled, digested and expanded according to previous Expansion Microscopy protocols.
4. Genomic loci are Fluorescent *in situ* hybridization (FISH) labeled after expansion.
5. Cells are imaged and the colocalization of immunofluorescent signal at each genomic loci is used as a quantitative measurement of histone mark levels for each locus.

SCEPTRE has been tested and validated for the labeling of histone H3K4me3 and H3K27me3 marks for specific genomic loci in h-TERT RPE1 cells. However, the method can be utilized for labeling other histone modifications or protein assemblies in the nucleus according to notes provided in "Before start". The method is also expected to work with cell lines or primary cells.

For more information about the method, see the below reference:

Woodworth, M.A., Ng, K.K.H., Halpern, A.R., Pease, N.A., Nguyen, P.H.B., Kueh, H.Y. and Vaughan, J.C. (2021) Multiplexed single-cell profiling of chromatin states at genomic loci by expansion microscopy. *Nucleic Acids Research*, 10.1093/nar/gkab423.

Scripts that can be used for image processing and analysis for SCEPTRE profiling can be found on github at:

https://github.com/marcwood13/SCEPTRE_pipeline

MATERIALS

Below is a list of materials for profiling of H3K4me3 and H3K27me3 histone marks at *GAPDH* loci in hTERT-RPE1 cells. For targeting of a different set of histone marks, use ChIP-grade antibodies specific to each histone mark of interest. For targeting a different genomic locus of interest, design DNA FISH oligonucleotide probes accordingly.

primary antibodies:

Rabbit anti-H3K4me3 (Active motif, 39159)

Mouse anti-H3K27me3 (Active Motif, 61017).

Secondary antibodies:

Donkey anti-rabbit (711-005-152) conjugated in-lab to Alexa Fluor 568

Donkey anti-mouse conjugated with Alexa Fluor 488 (715-545-150)

Enzymes:

proteinase K (Thermo Fisher Scientific, E00491)

RNase A (Thermo Fisher Scientific, EN0531)

alcohol oxidase (Sigma-Aldrich, A2404-1KU)

catalase (Sigma-Aldrich, C100)

Phusion Hot-start master mix (New England Biolabs, M0536L)

DNase I (New England Biolabs, M0303A)

Maxima H Minus RT Transcriptase (Thermo Fisher Scientific, EP0752).

Chemical reagents:

10× phosphate-buffered saline (PBS, Fisher Bioreagents, BP399-1)

32% paraformaldehyde aqueous solution (PFA, Electron Microscopy Sciences, RT15714)

4-(1,1,3,3-tetramethylbutyl)phenyl-polyethylene glycol (Triton X-100, Sigma-Aldrich, X100)

Bovine serum albumin (BSA, Rockland Immunochemicals Inc., BSA-50)

Alexa Fluor 568 NHS-ester (Thermo Fisher Scientific, A-20003)

methacrylic acid NHS-ester (MA-NHS, Sigma-Aldrich, 730300)

40% acrylamide aqueous solution (Bio-Rad Laboratories, 1610140)

2% bis-acrylamide aqueous solution (Bio-Rad Laboratories, 1610142)

97% sodium acrylate powder (Sigma-Aldrich, 408220)

ammonium persulfate (APS, Thermo Fisher Scientific, 17874)

tetramethylethylenediamine (TEMED, Thermo Fisher Scientific, 17919)

10× tris-acetate-EDTA (TAE, Fisher Bioreagents, BP2434-4)

guanidine hydrochloride powder (Sigma-Aldrich, G3272)

sodium azide (Sigma-Aldrich, S2002), poly-L-lysine (Sigma-Aldrich, P8920)

sodium bicarbonate (VWR, 470302), formamide (Fisher Chemical, F84-1)

20× saline sodium citrate (SSC, Sigma-Aldrich, S6639)

50% OmniPur Dextran Sulfate (EMD Millipore, 3730)

Tween 20 (Sigma-Aldrich, P9416), Hoechst 33258 (Sigma-Aldrich, B2883-25MG)

Tris Base (Fisher scientific, BP152-500)

methyl viologen dichloride hydrate (Sigma-Aldrich 856177)

L-ascorbic acid (Fisher scientific, A61-25)

Additional reagents:

Dulbecco's modified eagle medium (Gibco, 11995065)

penicillin and streptomycin (Gibco, 15140122)
nonessential amino acids (Gibco, 11140050)
fetal bovine serum (Gibco, 26140079)
0.25% trypsin-EDTA (Gibco, 25200056)

GAPDH probe set, adapter and reporters (see supplemental spreadsheet).

SAFETY WARNINGS

32% Paraformaldehyde is toxic and must be handled with care.

Formamide is flammable, toxic and a suspected teratogen. All steps involving warm or heated solutions with Formamide should be handled within a fume hood.

Guanidine hydrochloride is a known irritant. Dissolve in water within a fumehood.

Acrylamide and bis-acrylamide are irritants, carcinogenic and suspected teratogens. Handle with care.

Please refer to the Safety Data Sheet of each reagent for more information on any present safety and/or environmental hazards.

BEFORE START INSTRUCTIONS

Prepare the following solutions:

RPE1 Cell culture media (store at 4 °C for ~6 months)

- Dulbecco's modified eagle medium
- 100 units/mL of penicillin and streptomycin
- 1% (w/v) nonessential amino acids
- 10% (v/v) fetal bovine serum

Note: If using a different cell line, use the culture media appropriate for the cell line.

Permeabilization buffer (store at 4 °C for ~6 months)

- 1X PBS
- 0.1% (v/v) Triton-X 100

Blocking Buffer (store at 4 °C for ~6 months)

- 1X PBS
- 10% (w/v) BSA

- 3mM Sodium Azide

Primary solution:

- 2 µg/mL Rabbit X H3K4me3
- 2 µg/mL Mouse X H3K27me3
- 1X PBS
- 10% (w/v) BSA
- 3mM Sodium Azide

Note: Alternative Rabbit and Mouse antibodies can be use to target a different set of histone marks or nuclear protein structures.

Secondary solution:

- 3 µg/mL Donkey X Rabbit Alexa568
- 3 µg/mL Donkey X Mouse Alexa488
- 1X PBS
- 10% (w/v) BSA
- 3mM Sodium Azide

Note: If using an alternative set of primary antibodies, make sure the secondary antibodies label each primary antibody independently and can be visualized in separate fluorescence channels.

ExM monomer solution (store at 4 °C for 1 month)

- 1× PBS
- 2 M NaCl
- 2.5% (w/w) acrylamide
- 0.15% (w/w)
- N,N'-methylenebisacrylamide
- 8.625% (w/w) sodium acrylate

Digestion solution (store at 4 °C for ~6 months)

- 1× TAE
- 0.5% (v/v) Triton X-100
- 0.8 M guanidine HCl
- 8 units/mL proteinase K

Hybridization buffer (make fresh for each experiment)

- 2× SSC
- 50% (v/v) formamide
- 0.1% (v/v) Tween 20

Hybridization mixture (make fresh for each experiment)

- 2× SSC
- 50% formamide (v/v)
- 10% dextran sulfate (w/v)

- 0.1% (v/v) Tween 20
- 3 mM sodium azide
- 100nM GAPDH probe set or an oligonucleotide probe library capable of targeting a different locus of interest
- 110nM fluorescent oligonucleotide reporter and adapter

Note: If targeting an alternative locus of interest, use ~10-20 nM oligo probe library per kb of targeted genomic region, and 1–1.5× concentration of oligo reporters and adapters to oligo probe library.

Wash buffer (store at 4 °C for ~6 months)

- 2× SSC
- 0.1% (v/v) Tween 20

Cell Culture and Fixation

3d

1 Grow h-TERT RPE1 with **cell culture media** to ~80% confluency.

Step 1 includes a Step case.

Alt. cell line

step case

Alt. cell line

Instead of h-TERT RPE1 cells, an alternative adhesive cell line can be used for this protocol as long as:

1. DNA FISH probes are designed for targeting genomic regions in this cell line.
2. Nuclear expansion matches the expansion factor of the ExM gel recipe used. Proper expansion can be confirmed by comparing nuclear diameters of expanded vs. non-expanded cells.

Suspension cell lines may also work as long as the cell line can be attached to a coverslip before immunofluorescence steps.

2 Trypsinize cells with 0.25% trypsin-EDTA and seed at ~75,000 cells per well on top of round coverslips (no. 1.5, ~12 mm diameter) placed within 24-well culture plates.

3 Grow cells at 37 °C in 4% carbon dioxide humidified environment Overnight .

4 Briefly rinse cells with 250 µL of 1× PBS.

5 Fix cells with 250 μ L of cold EtOH:MeOH (1:1) for 00:06:00 at -20 $^{\circ}$ C . 6m

6 Wash three times with 250 μ L of 1 $\times$ PBS.

7 Store fixed cells in 250 μ L of 1 $\times$ PBS azide (1 $\times$ PBS with 3 mM sodium azide) at 4 $^{\circ}$ C .

Immunofluorescence

3d

8 Place fixed RPE1 cells in 250 μ L **permeabilization buffer** for 00:10:00 . 10m

9 Wash cells three times with 250 μ L of 1 $\times$ PBS.

10 Incubate in 250 μ L of **blocking solution** for 01:00:00 at Room temperature . 1h

11 Incubate in 250 μ L of **primary solution** Overnight ~18 hours at 4 $^{\circ}$ C .

12 Wash the sample with 250 μ L of **blocking solution** three times, 00:10:00 each time. 10m

13 Incubate in 250 μ L of **secondary solution** for 02:00:00 at Room temperature . 2h

14 Wash once for 00:10:00 with 250 μ L of **blocking solution**. 10m

15 Wash three times with 250 μ L of 1 $\times$ PBS.

16 Post-fix in 250 μ L of freshly made 4% PFA in 1 $\times$ PBS for 00:10:00 . 10m

17 Wash three times with 250 μ L of 1 $\times$ PBS.

18 Store immunofluorescently labeled cells in 250 μ L of 1 $\times$ PBS with 3mM sodium azide at 4 $^{\circ}$ C .

Gelation, Digestion and Expansion.

10m

19 Treat immunolabeled cells with 250 μ L of freshly prepared [M] 5 millimolar (mM) MA-NHS in 1 $\times$ PBS for 00:10:00 . 10m

20 Wash samples three times with 250 μ L of 1 $\times$ PBS.

21 Incubate in 250 μ L of **ExM monomer solution** for 00:10:00 . 10m

22 Remove coverslip with cell sample from 24-well plate and place, cells facing down, on to 50 μ L of **ExM monomer solution** supplemented with 0.15% (w/v) APS and 0.2% (w/w) TEMED.

23 Gel sample at Room temperature for 00:30:00 in a sealed container backfilled with nitrogen gas. 30m

24 Gently remove the cell-embedded hydrogel from the 12 mm coverslip.

25 Incubate cell-embedded hydrogel in 1 mL of **digestion solution** Overnight at 37 $^{\circ}$ C . 30m

26 Wash and expand by placing the digested sample in 50 mL of deionized water, replacing the water every 15-20 minutes for at least three times or until gel sample is $\sim$ 4 $\times$ the original size.

27 Store hydrogel sample at 4 °C in 2× SSC buffer.

*

II

DNA Fluorescence in situ hybridization

57m 30s

28 Slice a small (~3.5 mm × 3 mm × 2 mm, equivalent to 20 µL) piece of the hydrogel sample.

29 Incubate gel slice in 1 mL of **hybridization buffer** for 00:10:00 at Room temperature .

10m

30 Remove excess buffer and incubate gel slice in 500 µL of pre-heated **hybridization buffer** for 00:30:00 at 60 °C .

30m

30.1 Make 50 µL of **hybridization mixture** and pre-heat to 90 °C for 00:10:00 .

10m

31 Remove the excess buffer from the gel slice and add 50 µL of pre-heated hybridization mixture to the gel slice.

32 Denature sample at 90 °C for 00:02:30 .

2m 30s

33 Hybridize probes to sample Overnight at 42 °C . 5m

34 Remove excess buffer and wash gel slice by incubating in 500 μ L of preheated **wash buffer** 15m
for 00:15:00 at 60 °C .

35 Remove excess buffer and wash gel slice by incubating in 500 μ L of preheated **wash buffer** 15m
for 00:15:00 at 37 °C .

36 Remove excess buffer and wash gel slice by incubating in 500 μ L of **wash buffer** for 15m
00:15:00 at Room temperature .

37 Remove excess buffer and wash gel slice by incubating in 500 μ L of one tenth diluted **wash buffer** for 00:10:00 On ice . 10m

38 Store samples at 4 °C .

Expansion, imaging and analysis

10m

39 Remove excess buffer, then expand gel slice to ~4 $\times$ by adding and replacing 500 μ L of deionized water, twice every 00:10:00 at 4 °C . 10m

40 Place the expanded gel slice on a poli-lysine treated coverslip, cell-side facing down.

41 Image sample with a confocal microscope to obtain 3D image stacks of the nuclei of each cell.

42 Analyze fluorescence intensity of each histone modification at the individual loci to determine the activity state of each locus.