APR 24, 2021

OPEN ACCESS**DOI:**dx.doi.org/10.17504/protocols.io.bugwntxe

Protocol Citation: Oluwadamilola Osasona, Olumuyiwa Ariyo 2021. Mutational Analysis of Hepatitis B Virus Precore Region Molecular Variants in Ile-ife, Nigeria.

protocols.io<https://dx.doi.org/10.17504/protocols.io.bugwntxe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Apr 24, 2021**Last Modified:** Apr 24, 2021**PROTOCOL integer ID:**

49398

Mutational Analysis of Hepatitis B Virus Precore Region Molecular Variants in Ile-ife, Nigeria

Oluwadamilola Osasona¹, Olumuyiwa Ariyo¹¹Department of Biological Sciences, Redeemers University

Osasona Damilola

Tech. support email: osasona23@run.edu.ng

Oluwadamilola Osasona

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

Abstract

Introduction

Evolution of phenotypic diversity among viruses occurs as an escape mechanism against host immune pressure or drug selective pressure. Among HIV/HBV co-infected individuals, various HBV basal core promoter (BCP)/precore (PC) region molecular mutants had been reported with associated phenotypic defect in HBeAg production. The emergence of HBeAg negative variants of HBV in HIV co-infected individuals have profound implication on the diagnosis, management and prognosis of this subset of individuals. This includes delayed clearance of HBV, early development of adverse hepatic events such as liver cirrhosis and hepatocellular carcinoma . Currently, little is known about HBV BCP/PC region genomic heterogeneity in HIV/HBV co-infected patients in Nigeria. Therefore, this study was focussed on investigating evidence of precore/core region genomic variability among HIV/HBV co-infected patients in Nigeria.

Materials and methods

A total of 40 patients (20 HIV/HBV co-infected and 20 HBV mono-infected samples) were enrolled into the study and subsequently tested for HBsAg, HBeAg and HBeAb using specific Enzyme-Linked Immunosorbent Assay (ELISA). The BCP/PC genome regions (nucleotides 1653-1959) were amplified using a nested PCR assay and then subjected to BCP/PC mutational analysis in genome sites affecting HBeAg expression especially at the BCP transcriptional and PC Translational stop codon sites.

Results

Overall, 5(83.3%) of the six exploitable sequences after analysis showed various BCP/PC mutations. Only 1(16.6%) sequence from an HIV/HBV co-infected patient had the BCP transcriptional (double mutation; A1762T/G1764A) mutant. Analysis of the PC translational stop codon showed 4 (66.6%) having the G1896A mutants while 33.3% (2) had G1899A mutants.

Conclusion

This study has broadened the available evidence of BCP/PC region molecular mutants among HIV/HBV co-infected patients in Nigeria and assessed the difference of mutation prevalence in comparison with HBV mono-infected cohort . We therefore recommend that HIV/HBV co-infected patients be routinely screened for hepatitis B virus precore region mutants to improve their patient outcome.