MAY 29, 2019

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.2pygdpw](https://doi.org/10.17504/protocols.io.2pygdpw)

External link:
<https://doi.org/10.1371/journal.pone.0217560>

Protocol Citation: Matias Ruggieri, Carolina Berini, Nicolas Ducasa, Miroslav Malkovsky, Paul Fisch, Mirna Biglione 2019. n-PCR for 10 samples. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.2pygdpw>

MANUSCRIPT CITATION:

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: May 08, 2019

Last Modified: May 29, 2019

🌐 n-PCR for 10 samples

📖 PLOS One

Matias Ruggieri¹, Carolina Berini², Nicolas Ducasa², Miroslav Malkovsky³, Paul Fisch⁴, Mirna Biglione²

¹Institut für Klinische Pathologie, Universitätsklinikum Freiburg, German;

²Instituto de Investigaciones Biomédicas en Retrovirus y SIDA (INBIRS), CONICET-Universidad de Buenos Aires, Buenos Aires, Argentina;

³UW School of Medicine and Public Health, Madison, Wisconsin, USA;

⁴Institut für Klinische Pathologie, Universitätsklinikum Freiburg, Germany

Matias Ruggieri

ABSTRACT

HTLV-1 diagnosis using PCR is based on the amplification of the viral DNA sequences. This is usually conducted using a nested PCR which involves two rounds of DNA amplification reactions targeting the viral polymerase gene. In this case, we set up the conditions to analyze a pool of 10 samples in a single n-PCR detecting the presence of HTLV-1.

GUIDELINES

Pool of 10 samples in one nPCR

MATERIALS

MATERIALS

☒ GeneRuler 50 bp DNA Ladder **Thermo Fisher Scientific Catalog #SM0371**

☒ Taq DNA Polymerase (1000 U) **Qiagen Catalog #Cat No./ID: 201205**

☒ MT-2 cells **Contributed by users Catalog #237**

☒ Thermo Fisher Scientific Germany **Invitrogen - Thermo Fisher Catalog #10416014**

☒ 100 bp DNA Ladder **Invitrogen - Thermo Fisher Catalog #15628019**

☒ dNTP Set 100 mM Solutions **Thermo Fisher Scientific Catalog #R0182**

Keywords: nPCR, HTLV

- Gloves should be changed regularly, particularly after preparation and before the dispensing of master mixes into PCR plates or strips to avoid cross contamination.
- Be aware that virus genetic material is being use, so be carful.

BEFORE START INSTRUCTIONS

- To avoid contamination risk, DNA extraction, Pre PCR set-up (mastermix preparation), and post-PCR procedures should be performed at least in separate workplaces.
- Benches, work stations, centrifuges, vortexes, and pipettes must be cleaned with 70% ethanol before and after every PCR set-up.

1 Equipment and Materials

1. Pipettes
2. 10µl presterilized filter tips
3. 20µl presterilized filter tips
4. 200µl presterilized filter tips
5. 1000µl presterilized filter tips
6. Thermal Cycle (we used TProfessional TRIO)
7. Electrophoresis equipment

2 Reagents and Chemicals

- Taq DNA Polymerase kit (1000 U), by [Qiagen](#), Catalog #: [Cat No./ID: 201205](#)
- 1. Buffers (PCR Buffer and CoralLoad PCR Buffer)
- 2. MgCl₂
- 3. Q solution
- 4. Taq polymerase
- dNTPs
- Nuclease free H₂O
- Forward (F) and reverse (R) primers
- Agarose
- Markers: 50pb and/or 100pb (Thermo Fisher Scientific, *Germany*)

3 Preparation and storage of reagents

- Stock reagents for PCR (10X buffer, dNTPs, MgCl₂, Q solution and primers) and enzymes should be stored at -20°C.
- Stock reagents should be diluted into working concentrations and aliquoted at -20°C into separate tubes.
- Working aliquots of sample and control DNA as well as primary PCR (Nest-1) products must be kept at -20°C for long-term storage but can be kept in the fridge for daily use.

4 PCR summary protocol

- Primary PCR/ Nest-1 PCR should be done using PolEF and PolER primers.
- For each group of samples include positive control (DNA from MT2 cell line) and a negative control of PCR mix without a DNA template.
- Nest-2 PCR should be done using the primers PolIF and PolIR. Also the primers Actin F and Actin R are necessary to amplify human Actin gene.
- Once the PCR run is complete, remove tubes and briefly centrifuge to spin down products.
- Store PCR tubes in the fridge or freezer
- Run the samples on 2% Agarose gel electrophoresis using gel red nucleic acid stain.
- Score results as presence or absence of bands comparing with control bands.

5 PCR Protocol

- For each run, at least two controls (1 positive and 1 negative) should be included in addition to the samples.
- Prepare the PCR master mix calculating enough volume for number of the samples +8%, to account for pipetting loss and dead volume.
- Master Mix for Nest-1 PCR (**10X PCR buffer**, dNTPs, MgCl₂, primers, Taq polymerase and water)
- Master Mix for Nest-2 PCR (**10X CoralLoad PCR Buffer**, dNTPs, MgCl₂, primers, Taq polymerase and water)

6 Preparation of PCR (Nest-1 PCR) x 10

		Initial Conc entrat ion	Final Conc entrat ion	For 1 Tube
	PCR Buffer	10x	1 X	5 ul
	Q soluti on	5x	1X	10 ul
	MgI2	25m M	2,5m M	5 ul
	dNTP s	2mM	0,2m M	5 ul

PolEF	10mM	0,3mM	1,5 ul
PolER	10mM	0,3mM	1,5 ul
Taq polymerase		0,5 units	0,2 ul
DNA sample			2 ul (20 ng/μl) from each sample. 10 samples = 20 ul
Water			add to a final concentration of 50 ul

Preparation of PCR (Nest-2 PCR) x 10

	Initial Concentration	Final Concentration	For 1 Tube
Coral Load PCR Buffer	10x	1 X	2,5 ul
Q solution	5x	1X	5 ul
MgI2	25mM	2,5mM	2,5 ul
dNTPs	2mM	0,2mM	2,5 ul
PolIF	10mM	0,3mM	0,75 ul
PolIR	10mM	0,3mM	0,75 ul
Actin F	10mM	0,3mM	0,75 ul
Actin R	10mM	0,3mM	0,75 ul
Taq polymerase		0,5 units	0,1 ul

First reaction product			5 ul
Water			add to a final concentration of 25 ul

7 Primers

Round	Name	5'-3' oligonucleotide sequence
Nest-1 PCR	PolEF	TTTA GGTG CCCA AACT GGAG
Nest-1 PCR	PolER	GCAG GATA TTGG AAGC CTCA G3
Nest-2 PCR	PolIF	GCCC TCAT GCCA GTGT TTAC
Nest-2 PCR	PolIR	CCTG GAGA TGGG ATCA GGTA G
Nest-2 PCR	Actin F	ATCG AGCA CGGC ATCG TCAC CAAC
Nest-2 PCR	Actin R	GTTG AAGG TCTC AAAC ATGA TCTG

8 PCR Cycling conditions

Nest-1:

Pol EF + Pol ER

Temp eratur e (°C)	Time	
94	3 min	
94	10 sec	40 cycles
54	20 sec	
70	45 sec	
70	5 min	
4	Hold	

Nest-2:

Pol IF + Pol IR + Act F + Acr R

Temp eratur e (°C)	Time	
94	3 min	
94	10 sec	40 cycles
52	20 sec	
70	45 sec	
70	5 min	
4	Hold	