

Jul 23, 2019 Version 1

NEBNext dA-Tailing Module (NEB #E6053) V.1

DOI

dx.doi.org/10.17504/protocols.io.4t3gwqn



New England Biolabs¹, Menna Teffera¹

¹New England Biolabs

New England Biolabs (NEB)

Tech. support phone: +1(800)632-7799 email: info@neb.com



Isabel Gautreau

New England Biolabs

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.4t3gwqn

External link: <https://www.neb.com/products/e6053-nebnext-da-tailing-module#Product%20Information>

Protocol Citation: New England Biolabs, Menna Teffera 2019. NEBNext dA-Tailing Module (NEB #E6053). **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.4t3gwqn>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: June 27, 2019

Last Modified: June 22, 2023

Protocol Integer ID: 25179



Abstract

The NEBNext dA-Tailing Module has been optimized to efficiently incorporate a non-templated dAMP on the 3' end of a blunt DNA fragment (1). 3'-dA DNA tailing prevents concatamer formation during subsequent ligation steps.

DNA tailed with the NEBNext dA-Tailing module may be ligated to adaptors or cloning vectors with complementary dT overhangs. The NEBNext dA-Tailing Module is provided as a master mix to maximize efficiency and convenience in DNA sample preparation workflows.

The NEBNext dA-Tailing Module has been validated by sequencing with the Illumina Genome Analyzer II (Illumina, Inc.) in conjunction with the NEBNext End Repair Module, NEBNext Quick Ligation Module and Phusion® High-Fidelity PCR Master Mix.

For larger volume requirements, customized and bulk packaging is available by purchasing through the OEM/Bulks department at NEB. Please contact OEM@neb.com for further information.

Guidelines

Safe Stop Point: This is a point where you can safely stop the protocol and store the samples prior to proceeding to the next step in the protocol.

Caution: Signifies a step in the protocol that has two paths leading to the same point.

Color: A color listed before or after a reagent name indicates the cap color of the reagent to be added.

Materials

MATERIALS

⊗ NEBNext dA-Tailing Reaction Buffer **New England Biolabs Catalog #E6055**

⊗ Klenow Fragment (3→5 exo-) **New England Biolabs Catalog #E6054 in Kit E6053**

STEP MATERIALS

⊗ NEBNext dA-Tailing Reaction Buffer **New England Biolabs Catalog #E6055**

⊗ Klenow Fragment (3→5 exo-) **New England Biolabs Catalog #E6054 in Kit E6053**



Protocol materials

⊗ NEBNext dA-Tailing Reaction Buffer **New England Biolabs Catalog #E6055**

⊗ Klenow Fragment (3→5 exo-) **New England Biolabs Catalog #E6054 in Kit E6053**

⊗ NEBNext dA-Tailing Reaction Buffer **New England Biolabs Catalog #E6055**

⊗ Klenow Fragment (3→5 exo-) **New England Biolabs Catalog #E6054 in Kit E6053**

⊗ Klenow Fragment (3→5 exo-) **New England Biolabs Catalog #E6054 in Kit E6053**

⊗ NEBNext dA-Tailing Reaction Buffer **New England Biolabs Catalog #E6055**

Before start

Starting Material: 1–5 µg of of end repaired, blunt DNA (100–1000 bp).



dA Tailing

- 1 Mix the following components in a sterile microfuge tube:

| Component | Volume |
|------------------------------------|-----------------------------|
| End Repaired, Blunt DNA | variable |
| NEBNext dA-Tailing Reaction Buffer | 5 μ l |
| Klenow Fragment (3'→5' exo-) | 3 μ l |
| Sterile water | variable |
| Total Volume | 50 μl |

 NEBNext dA-Tailing Reaction Buffer **New England Biolabs Catalog #E6055**

 Klenow Fragment (3→5 exo-) **New England Biolabs Catalog #E6054 in Kit E6053**

- 2 Incubate in a thermal cycler for  00:30:00 at  37 °C .

30m

- 3 Purify DNA sample on one spin column or using AMPure XP beads.



Note

Note: for details on how this module is used in the NEBNext Library Prep for Illumina workflow, please see the manual for NEBNext DNA library Prep Master Mix Set for Illumina (NEB #E6040).