VERSION 1

JUL 22, 2019

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.4nqgvdw

External link:

<https://www.neb.com/-/media/catalog/datacards-or-manuals/manuale7595.pdf?rev=6d5ae698cf394a27864e788e6056e8d8>

Protocol Citation: New England Biolabs, Menna Teffera 2019. NEBNext Ultra II Ligation Module (NEB # E7595) for NEBNext Ultra II End Repair/dA Tailing Module (NEB #E7546). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.4nqgvdw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Jun 24, 2019

Last Modified: Jul 22, 2019

NEBNext Ultra II Ligation Module (NEB # E7595) for NEBNext Ultra II End Repair/dA Tailing Module (NEB #E7546) V.1

New England Biolabs¹, Menna Teffera¹

¹New England Biolabs

Menna Teffera

ABSTRACT

This module is part of the Ultra™ II workflow, and is optimized for use with the NEBNext® Ultra II End Repair/dA-Tailing Module (NEB #[E7546](#)), for Illumina®-compatible library construction.

The NEBNext Ultra II Ligation Module is optimized for use with the NEBNext Ultra II End Repair/dA-Tailing Module (NEB #[E7546](#)) or the NEBNext Ultra II FS DNA Module (NEB #[E7810](#)).

GUIDELINES

Safe Stop Point: This is a point where you can safely stop the protocol and store the samples prior to proceeding to the next step in the protocol.

Caution: Signifies a step in the protocol that has two paths leading to the same point.

Color: A color listed before or after a reagent name indicates the cap color of the reagent to be added.

Adaptor Dilution Guidelines

The appropriate adaptor dilution for your sample input and type may need to be optimized experimentally. The dilutions provided here are a general starting point.

Table 1.1: Adaptor Dilution

Input	Adapt or Dilution (Volume of adaptor: Total volume)	Working Adaptor Concentration
1 µg–101 ng	No Dilution	15 µM
100 ng–5 ng	10-Fold (1:10)	1.5 µM
less than 5 ng	25-Fold (1:25)	0.6 µM

MATERIALS

MATERIALS

Ligation Enhancer **New England Biolabs Catalog #E7374 in Kits E7370 or E7445**

NEBNext Adaptor for Illumina **New England Biolabs Catalog #E7337 in Kits E7335, E7500, E771**

NEBNext Ultra II Ligation Master Mix **New England Biolabs Catalog #E7648**

USER Enzyme (Multiplex Oligos for Illumina) **New England Biolabs Catalog #E7338**

STEP MATERIALS

USER Enzyme (Multiplex Oligos for Illumina) **New England Biolabs Catalog #E7338**

NEBNext Ultra II Ligation Master Mix **New England Biolabs Catalog #E7648**

NEBNext Ligation Enhancer **New England Biolabs Catalog #E7374**

NEBNext Adaptor for Illumina **New England Biolabs**

Materials that may be needed that are not included in this kit:

Tris-HCL Buffer

10 mM NaCl

BEFORE START INSTRUCTIONS

Starting Material:

500 pg–1 µg fragmented DNA that has been end repaired and dA-Tailed using the NEBNext End Repair/dA-Tailing Module ([NEB #E7546](#)).

Note

Caution: If DNA input is ≤ 100 ng, dilute the NEBNext Adaptor for Illumina in 10 mM Tris-HCl or 10 mM Tris-HCl with 10 mM NaCl as indicated in Table 1.1.

Ligation/End Prep

- 1 Add the following products directly to the End Prep Reaction Mixture:

Component	Volume
End Prep Reaction Mixture	60 µl

(red) NEBN ext Ultra II Ligati on Mast er Mix*	30 µl
(red) NEBN ext Ligati on Enhan cer	1 µl
(red) NEBN ext Adapt or for Illumi na**	2.5 µl
Total Volu me	93.5 µl

* Mix the Ultra II Ligation Master Mix by pipetting up and down several times prior to adding to the reaction.

** The NEBNext adaptor is provided in NEBNext Singleplex (NEB #E7350) or Multiplex (NEB #E7335, #E7500, #E7710, #E7730, #E7600, #E7535, and #E6609) Oligos for Illumina.

NEBNext Ultra II Ligation Master Mix **New England Biolabs Catalog #E7648**

NEBNext Ligation Enhancer **New England Biolabs Catalog #E7374**

NEBNext Adaptor for Illumina **New England Biolabs**

Note

Note: The Ligation Master Mix and Ligation Enhancer can be mixed ahead of time and is stable for at least 8 hours @ 4°C. We do not recommend premixing the Ligation Master Mix, Ligation Enhancer and adaptor prior to use in the Adaptor Ligation Step.

- 2 Set a 100 µl or 200 µl pipette to 80 µl and then pipette the entire volume up and down at least 10 times to mix thoroughly. Perform a quick spin to collect all liquid from the sides of the tube.

Note

Caution: The NEBNext Ultra II Ligation Master Mix is very viscous. Care should be taken to ensure adequate mixing of the ligation reaction, as incomplete mixing will result in reduced ligation efficiency. The presence of a small amount of bubbles will not interfere with performance.

- 3 Incubate at 20 °C for 00:15:00 in a thermocycler **with the heated lid off**.

15m

- 4 Add 3 µl of (red) USER™ Enzyme to the ligation mixture from Step 3.

USER Enzyme (Multiplex Oligos for Illumina) **New England Biolabs Catalog #E7338**

Note

Note: Steps 4 and 5 are only required for use with NEBNext Adaptors. USER enzyme can be found in the NEBNext Singleplex (NEB #E7350) or Multiplex (NEB #E7335, #E7500, #E7710, #E7730, #E7600 and #E6609) Oligos for Illumina.

- 5 Mix well and incubate at 37 °C for 00:15:00 with the heated lid set to ≥ 47 °C .

15m

- 6 DNA is now ready for size selection or cleanup.

Note

Note: Please see NEB #E7645 manual for recommended size selection/cleanup and PCR amplification protocols.

Note

Safe Stop Point: Samples can be stored overnight at –20°C.