FEB 27, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.x84fryw

Protocol Citation: Celeste M M. Karch, Rita Martinez, Jacob Marsh 2019. Neural aggregate formation.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.x84fryw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Feb 17, 2019

Last Modified: Feb 27, 2019

PROTOCOL integer ID:
 20476

Neural aggregate formation

In 1 collection

Celeste M M. Karch¹, Rita Martinez¹, Jacob Marsh¹

¹Washington University in St Louis

Neurodegeneration Method Development Community

Tech. support email: ndcn-help@chanzuckerberg.com

Celeste M M. Karch

Washington University in St Louis

ATTACHMENTS

[IPSC CORTICAL DIFFERENTIATION 022017.pdf](#)

GUIDELINES

This protocol is part of the [IPSC CORTICAL DIFFERENTIATION](#) collection.

This method should be performed using sterile technique.

MATERIALS

Please refer to the attached full manuscript for required materials.

SAFETY WARNINGS

Please refer to the SDS (Safety Data Sheet) for information about hazards, and to obtain advice on safety precautions.

- 1 Harvest iPSCs for neural aggregate formation when iPSCs have reached 75- 85% confluency. Aspirate medium and rinse with 2 mL of DPBS.

- 2 Add 1 mL of Accutase. Incubate at 37 °C for 00:05:00 . Gently tap plate to dislodge cells.
- 3 Dilute Accutase with 4 mL of DMEM/F12 medium and collect cell suspension in 15ml conical tube.
- 4 Centrifuge cells at 750 rpm for 00:03:00 . Then carefully aspirate medium from iPSC pellet.
- 5 Add 3 mL of neural induction medium to iPSC pellet. Using a hemacytometer, count iPSCs. Adjust volume of iPSC suspension to 450-650,000 cells/mL using neural induction medium supplemented Rock inhibitor (10 µM final).
- 6 Add 100 µL of iPSC suspension per well to a v-bottom 96-well plate.
- 7 Centrifuge plate at 750 rpm for 00:03:00 to sediment iPSC into spheres.
- 8 Incubate cells at 37 °C , 5% CO2 and 95% humidified chamber for 24:00:00 . After 24 hrs, carefully remove all medium from well and replace with 100 µL per well of Neural Induction Medium.

Note

Do not disturb or break apart spheres. The spheres are very delicate at this stage.

- 9 Incubate neurospheres in 96 well plate for 96:00:00 . Perform half volume medium changes daily (removed 50 μ L and replace with 50 μ L).