



Aug 15, 2019

🌐 Neuropathy Phentotyping Protocols - Basic Breeding to Harvest

DOI

dx.doi.org/10.17504/protocols.io.3jngkme



Eva Feldman¹

¹University of Michigan - Ann Arbor

Diabetic Complications Consortium
Tech. support email: rmcindoe@augusta.edu



Lili Liang

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.3jngkme>

External link: <https://www.diacomp.org/shared/document.aspx?id=54&docType=Protocol>

Protocol Citation: Eva Feldman 2019. Neuropathy Phentotyping Protocols - Basic Breeding to Harvest. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.3jngkme>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: May 31, 2019

Last Modified: August 15, 2019

Protocol Integer ID: 23886

Keywords: Neuropathy Phentotyping, Breeding, Diabetic Neuropathy, diabetic neuropathy at the whole animal, presence of diabetic neuropathy, development of diabetic neuropathy, diabetic neuropathy, neuropathy phentotyping protocol, detailed measures of neuropathy, assessments of neuropathy, diabetes, diabetic complication, neuropathy, duration of diabetes, initial degree of neuropathy, degree of diabetes, month period of diabetes, loss of nerve fiber, glycemic control, degree of glycemic control, glucose, tail blood glucose, glycated hemoglobin level, phenotyping of rodent, nerve fiber

Abstract

Summary:

Phenotyping of Rodents for the Presence of Diabetic Neuropathy

In man, the development of diabetic neuropathy is dependent on both the degree of glycemic control and the duration of diabetes. Diabetic neuropathy is a progressive disorder, with signs and symptoms that parallel the loss of nerve fibers over time. Consequently, assessments of neuropathy in mice are not performed at one time point, but are characterized at multiple time points during a 6 month period of diabetes. The degree of diabetes is evaluated in 2 ways: tail blood glucose measured following a 6 hour fast and glycated hemoglobin levels. The initial degree of neuropathy is screened using the methods discussed below. Detailed measures of neuropathy are employed when the initial screening instruments indicate a profound or unique phenotypic difference. This document contains protocols used by the DiaComp staff to examine and measure diabetic neuropathy at the whole animal, tissue and cellular levels.

Diabetic Complication:



Neuropathy

Troubleshooting

- 1 Basic breeding scheme is harem breeding: 2 females, 1 male per cage.
- 2 At least 3 breeding trios are active and are replaced every 6 months.
- 3 Tail biopsies are collect at 3-4 weeks of age for genotyping.
- 4 Fasting blood glucose tests are performed every 4 weeks.
- 5 Body weights are recorded every 4 weeks at same time as glucose test.
- 6 In animals not spontaneously diabetic, diabetes is induced by STZ injections at 8 weeks. 40mg/kg for 5 consecutive days.
- 7 Urine is collected at 12 and 24 weeks. (24 week time point is a 24 hour collection)