



Aug 16, 2019

🌐 Neuropathy Phentotyping Protocols - Dosages of Tranquilizers and Anesthetics for Mice, Rats, Hamsters/Gerbils

DOI

<https://dx.doi.org/10.17504/protocols.io.3jqgkmw>

Eva Feldman¹

¹University of Michigan - Ann Arbor

Diabetic Complications Consortium
Tech. support email: rmcindoe@augusta.edu



Lili Liang



Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.3jqgkmw>

External link: <https://www.diacomp.org/shared/document.aspx?id=54&docType=Protocol>

Protocol Citation: Eva Feldman 2019. Neuropathy Phentotyping Protocols - Dosages of Tranquilizers and Anesthetics for Mice, Rats, Hamsters/Gerbils. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.3jqgkmw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: May 31, 2019

Last Modified: August 16, 2019

Protocol Integer ID: 23888

Keywords: Diabetic Neuropathy , Neuropathy Phenotyping, Dosages of Tranquilizers and Anesthetics, diabetic neuropathy at the whole animal, presence of diabetic neuropathy, diabetic neuropathy, development of diabetic neuropathy, detailed measures of neuropathy, assessments of neuropathy, neuropathy phenotyping protocol, neuropathy, diabetes, duration of diabetes, initial degree of neuropathy, diabetic complication, loss of nerve fiber, degree of diabetes, glycemic control, month period of diabetes, anesthetics for mice, degree of glycemic control, tail blood glucose, glycated hemoglobin level, nerve fiber, phenotyping of rodent

Abstract

Summary:

Phenotyping of Rodents for the Presence of Diabetic Neuropathy

In man, the development of diabetic neuropathy is dependent on both the degree of glycemic control and the duration of diabetes. Diabetic neuropathy is a progressive disorder, with signs and symptoms that parallel the loss of nerve fibers over time. Consequently, assessments of neuropathy in mice are not performed at one time point, but are characterized at multiple time points during a 6 month period of diabetes. The degree of diabetes is evaluated in 2 ways: tail blood glucose measured following a 6 hour fast and glycated hemoglobin levels. The initial degree of neuropathy is screened using the methods discussed below. Detailed measures of neuropathy are employed when the initial screening instruments indicate a profound or unique phenotypic difference. This document contains protocols used by the DiaComp staff to examine and measure diabetic neuropathy at the whole animal, tissue and cellular levels.

Diabetic Complication:



Neuropathy

Materials

Dosages of Tranquilizers and Anesthetics for Mice, Rats, Hamsters/Gerbils

Tranquilizer and Anesthetic	Mouse mg/kg (route)	Rat mg/kg (route)	Hamster/Gerbil mg/kg (route)
Promazine (Sparine)	4 (IM,SC)	0.5-1 (IM)	0.5-1 (IM)
Acepromazine (PromAce)	0.75 (IM,SC)	0.75 (IM)	
Chlorpromazine (Thorazine)	5-10 (IM)	1-2 (IM)	0.5 (IM)
Xylazine (Rompun)	2.5 (IM)	2.5 (IM)	10 (IM)
Diazepam (Valium)	5.0 (IP)	2.5 (IP)	5.0 (IP)
Droperidol and Fentanyl (Innovar-Vet)	0.2-0.5 ml/kg (IM)	0.1-0.5 ml/kg (IM)	contraindicated
Ketamine (Ketastet)	50-80 (IM)a	20-60 (IM)a	40-150 (IM)a 100-200 (IP)a
Pentobarbital (Diabutal, Nembutal)	35-70 (IV) 60-90 (IP)	25-40 (IV) 35-50 (IP)	30 (IV) 50-90 (IP)
Thiopental (Pentothal)	25-50 (IV) 50 (IP)	20 (IV) 20-40 (IP)	20 (IV) 40 (IP)
Thiamylal (Surital, Bio-tal)	25-50 (IV)	20 (IV) 20-25 (IP)	

a - Recommend to combine with Xylazine or Acepromazine.

