AUG 16, 2019

Neuropathy Phenotyping Protocols - Dosages of Tranquilizers and Anesthetics for Mice, Rats, Hamsters/Gerbils

Eva Feldman¹

¹University of Michigan - Ann Arbor

Diabetic Complications Consortium

Tech. support email: rmcindoe@augusta.edu

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.3jqgkmw

External link:
<https://www.diacomp.org/shared/document.aspx?id=54&docType=Protocol>

Protocol Citation: Eva Feldman 2019. Neuropathy Phenotyping Protocols - Dosages of Tranquilizers and Anesthetics for Mice, Rats, Hamsters/Gerbils.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.3jqgkmw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 31, 2019

Last Modified: Aug 16, 2019

Lili Liang

ABSTRACT

Summary:

Phenotyping of Rodents for the Presence of Diabetic Neuropathy

In man, the development of diabetic neuropathy is dependent on both the degree of glycemic control and the duration of diabetes. Diabetic neuropathy is a progressive disorder, with signs and symptoms that parallel the loss of nerve fibers over time. Consequently, assessments of neuropathy in mice are not performed at one time point, but are characterized at multiple time points during a 6 month period of diabetes. The degree of diabetes is evaluated in 2 ways: tail blood glucose measured following a 6 hour fast and glycated hemoglobin levels. The initial degree of neuropathy is screened using the methods discussed below. Detailed measures of neuropathy are employed when the initial screening instruments indicate a profound or unique phenotypic difference. This document contains protocols used by the DiaComp staff to examine and measure diabetic neuropathy at the whole animal, tissue and cellular levels.

Diabetic Complication:

Neuropathy

Keywords: Diabetic
Neuropathy , Neuropathy
Phentotyping, Dosages of
Tranquilizers and Anesthetics

Dosages of Tranquilizers and Anesthetics for Mice, Rats, Hamsters/Gerbils

Tranquilizer and Anesthetic	Mouse mg/kg (route)	Rat mg/kg (route)	Hamster/Gerbil mg/kg (route)
Promazine (Sparine)	4 (IM,SC)	0.5-1 (IM)	0.5-1 (IM)
Acepromazine (PromAce)	0.75 (IM,SC)	0.75 (IM)	
Chlorpromazine (Thorazine)	5-10 (IM)	1-2 (IM)	0.5 (IM)
Xylazine (Rompun)	2.5 (IM)	2.5 (IM)	10 (IM)
Diazepam (Valium)	5.0 (IP)	2.5 (IP)	5.0 (IP)
Droperidol and Fentanyl (Innovar-Vet)	0.2-0.5 ml/kg (IM)	0.1-0.5 ml/kg (IM)	contraindicated
Ketamine (Ketastet)	50-80 (IM)a	20-60 (IM)a	40-150 (IM)a 100-200 (IP)a
Pentobarbital (Diabutal, Nembutal)	35-70 (IV) 60-90 (IP)	25-40 (IV) 35-50 (IP)	30 (IV) 50-90 (IP)
Thiopental (Pentothal)	25-50 (IV) 50 (IP)	20 (IV) 20-40 (IP)	20 (IV) 40 (IP)
Thiamylal (Surital, Bio-tal)	25-50 (IV)	20 (IV) 20-25 (IP)	

a - Recommend to combine with Xylazine or Acepromazine.