AUG 16, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.3jwgkpe

External link:
<https://www.diacomp.org/shared/document.aspx?id=54&docType=Protocol>

Protocol Citation: Eva Feldman 2019. Neuropathy Phenotyping Protocols - Immunofluorescence Method for 8DG Localization.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.3jwgkpe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: May 31, 2019

Last Modified: Aug 16, 2019

PROTOCOL integer ID:
 23894

Neuropathy Phenotyping Protocols - Immunofluorescence Method for 8DG Localization

Eva Feldman¹

¹University of Michigan - Ann Arbor

Diabetic Complications Consortium
 Tech. support email: rmcindoe@augusta.edu

Lili Liang

ABSTRACT

Summary:

Phenotyping of Rodents for the Presence of Diabetic Neuropathy

In man, the development of diabetic neuropathy is dependent on both the degree of glycemic control and the duration of diabetes. Diabetic neuropathy is a progressive disorder, with signs and symptoms that parallel the loss of nerve fibers over time. Consequently, assessments of neuropathy in mice are not performed at one time point, but are characterized at multiple time points during a 6 month period of diabetes. The degree of diabetes is evaluated in 2 ways: tail blood glucose measured following a 6 hour fast and glycated hemoglobin levels. The initial degree of neuropathy is screened using the methods discussed below. Detailed measures of neuropathy are employed when the initial screening instruments indicate a profound or unique phenotypic difference. This document contains protocols used by the DiaComp staff to examine and measure diabetic neuropathy at the whole animal, tissue and cellular levels.

Diabetic Complication:

Neuropathy

Reference:

Yarborough et al, Cancer Res. 56:683-688, 1996

MATERIALS

Reagents:

- RNase
- Proteinase K
- HCl
- Ethos, 50, 70, 95, 100%
- Hemo-De
- PBS
- Tris (10 mM, pH 7.5, 1 mM EDTA, 0.4 M NaCl)

Counterstains:

- Bis-benzimide

Other Chemicals:

- Normal serum
- Non-fat dry milk
- Bleach
- H₂O₂

Mounting Media:

- Prolong
- Gelmount

Glassware:

- Square Wheaton dishes
- Slide racks and handles
- Coplin jars
- Beakers
- Microfuge tubes

- 1 Deparaffinize slides by soaking in Hemo-D overnight in rocker oven. If using cryosections, thaw on warm plate 10 min, ring with PAP pen.
- 2 Re-hydrate through EtOH 100% -50% dH₂O.
- 3 Equilibrate sections in Tris. (10 mM, pH 7.5, 1 mM EDTA, 0.4 M NaCl)

- 4 Incubate with RNase in Tris, 100 µg/ml, 37°C, 1 hour.
- 5 Rinse 2 X 10 min Tris.
- 6 Incubate with Proteinase K 10 µg/ml Tris 22°C, 7 min.
- 7 Rinse 2 X 10 min Tris.
- 8 Incubate in 4 N HCl 22°C, 7 min.
- 9 Rinse 2 X 10 min Tris.
- 10 Rinse 2 X 10 min PBS.
- 11 Begin normal IHC procedure with the 8DG antibody.