VERSION 2

AUG 12, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.6f2hbqe

Protocol Citation: Khairil Asnan, Mohd. Sadzan Asram bin Aliming, Aprialdy Idrus, Diana Eka Pratiwi 2019. New Iron Extracting Method from Cattle's Blood for Iron Concentration Analysis.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.6f2hbqe>

MANUSCRIPT CITATION:

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol in our group and it is working. We will develop this protocol for better result

Created: Aug 12, 2019

🌐 New Iron Extracting Method from Cattle's Blood for Iron Concentration Analysis V.2

Khairil Asnan¹, Mohd. Sadzan Asram bin Aliming²,
 Aprialdy Idrus², Diana Eka Pratiwi²

¹Biology Department, Universitas Negeri Makassar;

²Chemistry Department, Universitas Negeri Makassar

Khairil Asnan

ABSTRACT

Cattle's blood component is compressed using centrifuge (5000 rpm for 10 minutes), then supernatant (erythrocytes) collected. Erythrocytes with a ratio of 3:1 (serum:supernatant), treated with initial mixing with NaOH (0.5 M) with an initial ratio 1:1 (v/v) and let it sit for 30 seconds then mixed it by centrifugation about 10 rpm for 30 seconds. Sample then treated by mixing oleic acid (2:1 v/v). The sample then dehydrated by heat about 121°C for a week. Calculated total iron mass was 240000 µg/100 ml whole blood sample (about 14.40% content of the whole sample). Sample purified by furnace using high temperature about 800°C for 2 hours and increased the iron concentration up to 46.30% (m/m%).

MATERIALS

MATERIALS

☒ Sodium Hydroxide Merck MilliporeSigma (Sigma-Aldrich)

STEP MATERIALS

☒ Sodium Hydroxide Merck MilliporeSigma (Sigma-Aldrich)

we also used oleic acid as chelating agent

Last Modified: Aug 12, 2019

PROTOCOL integer ID:
26842

Keywords: Physiology,
Biology, Bioanalysis,
Biosynthesis, Iron-extracting
method

Sample preparation

1

Compress the whole blood sample from cattle. 6 mL 37 °C 00:10:00

Equipment

Mini-centrifuge

NAME

Centrifuge

TYPE

Fisher

BRAND

S67601B

SKU

<https://www.fishersci.com/shop/products/fisherbrand-standard-mini-centrifuge-standard-mini-centrifuge/s67601b>

LINK

Any standard mini centrifuge with adapters for different tube sizes will suffice SPECIFICATIONS

5000 rpm

2

Mix 2 mL blood sample sample with 2 mL NaOH 37 °C 00:00:30

[M] 0.5 Molarity (M)

Sodium Hydroxide **Merck MilliporeSigma (Sigma-Aldrich)**

Expected result

dark green solution with strong odor

🌀 10 rpm , after we rest the mixed solution for 30 seconds

- 3 Chelate reaction by adding 🧪 4 mL Oleic acid into the previous mixed sample solution then let it sit for 🕒 00:00:30 . After that, mixed them by 🌀 10 rpm for 🕒 00:00:30 .

Expected result

Equipment

Centrifuge

NAME

Benchtop Centrifuge

TYPE

Eppendorf

BRAND

5405000441

SKU

<https://online-shop.eppendorf.us/US-en/Centrifugation-44533/Centrifuges-44534/Centrifuge-5425-PF-243560.html>

LINK

Any benchtop centrifuge will suffice

SPECIFICATIONS