JUL 29, 2019

Nile Red Staining of Drosophila Larval Tissues

Forked from [Nile Red Staining of Drosophila Larval Tissues](#)

Elizabeth Allen¹

¹University of Massachusetts Medical School

Jiajun Li

ABSTRACT

This protocol is used to stain late larval *Drosophila* lipid droplets in fat bodies and intestines with Nile Red, which emits fluorescence in the 552/636 nm range.

GUIDELINES

In brief, dissect animals in ice cold phosphate-buffered saline (PBS). Keep tissues in PBS on ice while obtaining your desired sample size. Fix tissues in 4% PFA (diluted in PBS with 0.1% Triton X-100), for 20 minutes, wash 3x in PBS, and stain tissues light-protected at room temperature for 1 hour.

* Do not use any serums for this protocol because Nile Red will instead be drawn away from your tissues and into serum.

MATERIALS

MATERIALS

- ⊗ 1X PBS (Phosphate-buffered saline) **Contributed by users**
- ⊗ Triton-X100 **Contributed by users**
- ⊗ Paraformaldehyde Powder (PFA) **Contributed by users Catalog #P6148**
- ⊗ Nile Red **Sigma Aldrich Catalog #N3013 SIGMA**
- ⊗ Acetone solution **Sigma Aldrich Catalog #48358 SUPELCO**

STEP MATERIALS

- ⊗ 1X PBS (Phosphate-buffered saline) **Contributed by users**
- ⊗ 1X PBS (Phosphate-buffered saline) **Contributed by users**
- ⊗ 1X PBS (Phosphate-buffered saline) **Contributed by users**

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.5x7g7rn

Protocol Citation: Elizabeth Allen 2019. Nile Red Staining of Drosophila Larval Tissues. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.5x7g7rn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jul 29, 2019

Last Modified: Jul 29, 2019

PROTOCOL integer ID:
 26335

BEFORE START INSTRUCTIONS

1. Prepare Nile Red (Sigma-Aldrich) in acetone (1000 ug/mL).
2. Dissect tissues in ice cold PBS.
3. Fix tissues in 4% paraformaldehyde (formalin) diluted in PBS with 0.1% Triton X-100.
4. Wash tissues 3x in PBS.

- 1 Dissect tissues in ice cold PBS, keeping samples on ice until required sample size is obtained.

1X PBS (Phosphate-buffered saline) **Contributed by users**

- 2 Fix tissues in 4% PFA for 20 minutes.

00:20:00

- 3 Wash tissues 3x in PBS.

1X PBS (Phosphate-buffered saline) **Contributed by users**

- 4 Stain samples in Nile Red at 0.5 ug/mL diluted in PBS for 1 hour.

01:00:00

Protocol

NAME

Nile Red prepared in acetone

CREATED BY

Elizabeth Allen

PREVIEW

- 4.1 Prepare a concentrated working solution of Nile Red in acetone at 1000 ug/mL.

Nile Red **Sigma Aldrich Catalog #N3013 SIGMA**

- 4.2 Store concentrated Nile Red solution at 4°C in the dark for up to 3 months.

4.3 Use Nile Red/acetone concentrate diluted in PBS at a concentration of 0.5 ug/mL.

*adjust duration of staining according to the tissue type, and stain at room temperature in the dark.

5 Wash tissues 3x in PBS.

1X PBS (Phosphate-buffered saline) **Contributed by users**

6 Carefully replace the PBS with mounting medium before transferring samples to slides and imaging.

* Image immediately, or temporarily store slides at 4°C.

Protocol

NAME

Mounting Media for Immunohistochemistry - Drosophila

CREATED BY

Sonia M Hall

PREVIEW

6.1 90% glycerol
10% 1M Tris-base pH 8.0
0.5% n-propyl-gallate