SEP 11, 2019

Nodule cleaning

In 1 collection

Ming-Dao Chia

ABSTRACT

Adapted for high throughput of small samples for sequencing.

Based on doi:10.1371/journal.pone.0047677

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.s4xegxn

Protocol Citation: Ming-Dao Chia: Nodule cleaning.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.s4xegxn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

These are protocols used in the Honours thesis of the same title:

<https://doi.org/10.25911/5d6cf6fb99437>

Created: Aug 29, 2018

Last Modified: Sep 11, 2019

PROTOCOL integer ID:

15223

- 1 Cleanly cut nodules from plant roots and drop into a 2ml Eppendorf tube per plant.

- 2 Add 1ml 100% EtOH to each tube
- 3 Incubate for 5 seconds
- 4 Remove as much liquid as possible without removing loose nodules
- 5 Add 1ml 6% sodium hypochlorite
- 6 Remove as much liquid as possible
- 7 Add 2ml sterile water
- 8 Remove as much liquid as possible

- 9 Close tube, move to sequencing/culturing