JAN 21, 2021

nPCR measurement method

PLOS One

lyzhaojh¹

¹, Zhejiang People's Hospital, Yi-Wen Li

lyzhaojh

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bp94mr8w

External link:

<https://doi.org/10.1371/journal.pone.0245370>

Document Citation: lyzhaojh 2021. nPCR measurement method. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.bp94mr8w>

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Dec 01, 2020

Last Modified: Jan 21, 2021

DOCUMENT integer ID:

45084

1. Clinically, we will encounter only limited data. For example, we only have data on urea nitrogen before hemodialysis and KT/V for single dialysis. How to calculate the normalized protein catabolic rate (nPCR)?
2. We introduce a simple formula, as followed. $nPCR(g/kg/d) = C_0/[a+b*KT/V+c/(KT/V)]+0.168$, where C_0 indicates pre-dialysis blood nitrogen in md/dl; a,b,c has different coefficients depending on the time of the dialysis schedule.
3. In our blood purification center, patients are dialyzed three times a week, either on Mondays, Wednesdays, Fridays or either Tuesdays, Thursdays, Saturdays, so that the time between the first dialysis sessions at the beginning of

the week is longer.

4. Therefore, the following formula is used: beginning-of-week:

$$nPCR(g/kg/d) = C_0 / [36.3 + 5.48 \cdot KT/V + 53.5 / (KT/V)] + 0.168$$

5. As 1 mmol/l urea nitrogen equal 2.802mg/dl, $nPCR(g/kg/d) = 2.802 \cdot BUN / [36.3 + 5.48 \cdot KT/V + 53.5 / (KT/V)] + 0.168$ (urea nitrogen is in mmol/l).