SEP 22, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bkacksaw

Protocol Citation: Danh Truong, Salah-Eddine Lamhamedi-Cherradi, Joseph A. Ludwig 2020. Nuclei Isolation for Single-Nuclei RNA sequencing. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bkacksaw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 25, 2020

Last Modified: Sep 22, 2020

PROTOCOL integer ID:
 40996

Keywords: single-cell, single-nuclei, nuclei isolation, fatty tissue, adipose tissue, liposarcoma, sarcoma, PDX

Nuclei Isolation for Single-Nuclei RNA sequencing

Danh Truong¹, Salah-Eddine Lamhamedi-Cherradi¹, Joseph A. Ludwig¹

¹MD Anderson Cancer Center

Human Cell Atlas Method Development Community

Ludwig Lab

Danh Truong
 MD Anderson Cancer Center

ABSTRACT

We developed this protocol while trying to isolate single-nuclei from frozen liposarcoma tissue. We tested this protocol on a variety of PDX's. For fattier tissues, we use a modified protocol based on a different kit. Thanks to Dr. Luciano Martelotto for his assistance and tips on the first section.

This protocol is based on prior work: ['Frankenstein' protocol](#), [Nuclei Isolation Kit: Nuclei EZ Prep](#), and [Minute™ Total Protein Extraction Kit for Adipose Tissues/Cultured Adipocytes](#).

GUIDELINES

Refer to [Isolation of Nuclei for Single Cell RNA Sequencing](#) from 10x genomics for additional information.

For filtering the nuclei, we don't use Flowmi filters as we find that some of the tips do not fit properly into the filters and I lost some samples due to this. I prefer to use the pluriStrainer Mini.

For nuclei sequencing, since nuclei have lower RNA content compared to whole cell, protocols could increase the number of cDNA amplification cycles by 1 or 2 cycles, as suggested by 10x genomics.

MATERIALS

MATERIALS

- ✂ RNAlater **Thermo Fisher Scientific Catalog #AM7020**
- ✂ DAPI **Invitrogen - Thermo Fisher Catalog #D3571**
- ✂ Protein LoBind Tubes, 1.5 mL **Eppendorf Catalog #0030108116**
- ✂ RNase-Free Disposable Pellet Pestles, With tube **Thermo Fisher Catalog #12141368**
- ✂ RNase Inhibitor **Thermo Fisher Catalog #N8080119**
- ✂ Nuclei Isolation Kit: Nuclei EZ Prep **Sigma Aldrich Catalog #NUC-101**
- ✂ PluriStrainer Mini 40µm **pluriSelect Catalog #43-10040-60**
- ✂ Minute Nuclei and Cytosol Isolation Kit for Adipose Tissues/Cultured Adipocytes **Fisher Scientific Catalog #AN-029**
- ✂ BioVortexer Mixer **Fisher Scientific Catalog #50-153-2378**
- ✂ pluriStrainer Mini 70 UM **Fisher Scientific Catalog #NC1444112**

STEP MATERIALS

- ✂ pluriStrainer 40 µm **pluriSelect Catalog #43-50040**
- ✂ Nuclei EZ lysis buffer **Merck MilliporeSigma (Sigma-Aldrich) Catalog #EZ PREP NUC-101**
- ✂ Minute Nuclei and Cytosol Isolation Kit for Adipose Tissues/Cultured Adipocytes **Fisher Scientific Catalog #AN-029**
- ✂ BioVortexer Mixer **Fisher Scientific Catalog #50-153-2378**
- ✂ RNase-Free Disposable Pellet Pestles, With tube **Thermo Fisher Catalog #12141368**
- ✂ BioVortexer Mixer **Fisher Scientific Catalog #50-153-2378**
- ✂ RNase-Free Disposable Pellet Pestles, With tube **Thermo Fisher Catalog #12141368**
- ✂ pluriStrainer Mini 70 UM **Fisher Scientific Catalog #NC1444112**

Consumables:

- Scalpels
- 1.5 mL tubes
- 15 mL centrifuge tubes
- 50 mL centrifuge tubes
- PBS, sterile

BEFORE START INSTRUCTIONS

All solutions should be kept at 4 °C or on On ice prior to use.

Turn on benchtop centrifuge and keep at 4 °C

Prepare Nuclei Wash and Resuspension Buffer (4 °C)

- 1X PBS with 1.0% BSA and 0.2U/μl RNase Inhibitor

Nuclei Isolation for Fatty tissues

- use a thermometer to make sure freezer is about -20 °C
- Chill plastic consumables in ice before use

Nuclei Isolation (most tissues)

25m

- 1 This protocol should work for most tissues. If the tissue that you are working is fatty or contains fat, including adipose or brain, **skip to Nuclei Isolation for Fatty tissues section.**

- 2 Use a scalpel and mince tissue on ice. Mincing tissue will improve nuclei isolation efficiency.

- 3 Add 500 μL of EZ Lysis Buffer to tissue in a 1.5 mL microcentrifuge tube.

Nuclei EZ lysis buffer **Merck MilliporeSigma (Sigma-Aldrich) Catalog #EZ PREP NUC-101**

- 4 Homogenize tissue using plastic microcentrifuge pestles. A Chemglass Life Sciences Supplier BioVortexer Mixer [Fisher Scientific: 50-153-2378] can be used. This should be done in ice but you can take it out for visual inspection. The timing will be based on tissue type and size.

45s

BioVortexer Mixer **Fisher Scientific Catalog #50-153-2378**

RNase-Free Disposable Pellet Pestles, With tube **Thermo Fisher Catalog #12141368**

- 5 Add 1 mL of EZ Lysis Buffer to the 1.5 mL microcentrifuge tube and incubate On ice for

5m

00:05:00 . This time can be modified depending on your tissue type and size.

- 6 Filter the solution using pluriStrainer Mini 70 μm [Fisher Scientific: NC1444112] into a new 1.5 mL microcentrifuge tube.

pluriStrainer Mini 70 UM **Fisher Scientific Catalog #NC1444112**

- 7 Centrifuge the solution at 500 x g, 4°C, 00:05:00 .

5m

- 8 Add 1 mL of EZ Lysis Buffer to the 1.5 mL microcentrifuge tube and incubate On ice for 00:05:00 .

- 9 Centrifuge the solution at 500 x g, 4°C, 00:05:00 .

- 10 Remove the supernatant while being careful not to disturb the pellet. If you can not see the pellet, it is advisable to leave behind 50 μL .

Washing

15m

- 11 Carefully add 0.5 mL of the **Nuclei Wash and Resuspension Buffer** and incubate for 00:05:00 without resuspending.

- 12 Add 0.5 mL of the **Nuclei Wash and Resuspension Buffer** and resuspend the nuclei.

- 13 Centrifuge the solution at 500 x g, 4°C, 00:05:00 . 5m
- 14 Aspirate the supernatant while being careful not to disturb the pellet and leave behind 50 µL .
- 15 Add 1 mL of **Nuclei Wash and Resuspension Buffer**.
- 16 Centrifuge the solution at 500 x g, 4°C, 00:05:00 . 5m
- 17 Aspirate the supernatant and resuspend the nuclei in 500 µL of **Nuclei Wash and Resuspension Buffer** supplemented with 10 µg/µL DAPI.
- 18 Filter nuclei with a pluriStrainer Mini 40 µm [Fisher Scientific: NC1469671].
pluriStrainer 40 µm **pluriSelect Catalog #43-50040**

Visual Inspection and Sorting

1h

- 19 Visually inspect nuclei integrity under a microscope and count the number of nuclei with a cell counter. This is important to check before continuing to see if the protocol was successful.

Image of isolated nuclei. There is significant debris prior to sorting.

20

Nuclei should be sorted as there will be significant debris after dissociation.

45m

21 Nuclei can be visualized after sort to confirm success.

Image of isolated nuclei after sorting.

Image of isolated nuclei after sorting.

- 22 After sorting, adjust concentration of nuclei to 700 nuclei/ μ L if needed. Continue with 10x Genomics Single Cell Protocol.

Nuclei Isolation for Fatty tissues

10m

- 23 For fatty tissues, including brain and adipose, the lipids can interfere with the collection. This is a different protocol that can assist in enhancing nuclei isolation efficiency. We will use a modified protocol from the Minute Nuclei and Cytosol Isolation Kit for Adipose Tissues/Cultured Adipocytes kit.

Minute Nuclei and Cytosol Isolation Kit for Adipose Tissues/Cultured Adipocytes **Fisher Scientific Catalog #AN-029**

- 24 Tissue weight should range from 120 mg to 150 mg. Mince tissue using a scalpel and place inside a 1.5 mL microcentrifuge tube.

- 25 Add 600 μ L of **N/C buffer** to tissue in a 1.5 mL microcentrifuge tube.

- 26 Homogenize tissue using plastic microcentrifuge pestles. A Chemglass Life Sciences Supplier BioVortexer Mixer [Fisher Scientific: 50-153-2378] can be used. This should be done in ice but you can take it out for visual inspection. The timing will be based on tissue type and size.

BioVortexer Mixer **Fisher Scientific Catalog #50-153-2378**

RNase-Free Disposable Pellet Pestles, With tube **Thermo Fisher Catalog #12141368**

Filter Nuclei and Lipids

30m

- 27 Place the filter cartridge that comes with the kit onto the collection tube. Pour the homogenized

tissue and liquid into the filter. Pipette any remaining liquid into the filter.

28 Incubate the filter cartridge with the cap open in a freezer at -20 °C for 00:20:00 20m

28.1 The timing can be optimized if you do not have a thermometer. Add 0.5 mL of ddH₂O in a 1.5 mL microcentrifuge tube and incubate in the freezer until the liquid freezes. This is the time that can be used.

29 Place the tube with cap open in a microcentrifuge. Centrifuge at 500 x g, 4°C, 00:03:00 . Increase to 1000 x g, 4°C, 00:01:00 if there is some liquid retention. You will see fat stuck on top of filter. The supernatant below will contain the nuclei.

30 Discard the filter and close the microcentrifuge tube. Vortex briefly to mix up the nuclei. Make sure you know where the pellet will end up based on the placement inside the centrifuge. The nuclei pellet will be white or almost invisible. Centrifuge at 1000 x g, 4°C, 00:04:00 .

31 The supernatant can be saved if needed. It is the cytosolic fraction. I did not have a visible nuclei pellet but I marked where the pellet would be. Carefully aspirate the liquid and leave some behind if needed. Resuspend nuclei in 50 µL **Nuclei Wash and Resuspension Buffer**.

32 Transfer nuclei to a clean microcentrifuge tube to avoid lipid contamination leftover on the tube walls.

33 The nuclei can be visualized using trypan blue. You will notice lipid contamination still, which can be removed using FACS.

Isolated nuclei from fatty tissue.

Isolated nuclei from fatty tissue.

34

While both protocols will result in many lipid droplets in addition to the nuclei, the fatty tissue protocol will have a higher nuclei count overall.

Left graph shows results from the standard protocol. Right graph shows the results from the fatty tissue protocol.

35

go to step #22 for sorting prior to 10x Genomics Single Cell protocol.

Cryopreservation

1h 30m

36

Add 900 μL RNALater to 100 μL nuclei in **Nuclei Wash and Resuspension Buffer**, incubate at 4 $^{\circ}\text{C}$ for 01:00:00 then transfer to -80 $^{\circ}\text{C}$

1h

37

To remove RNALater, centrifuge nuclei at 4000 x g, 4 $^{\circ}\text{C}$, 00:10:00. Aspirate RNALater and resuspend in **Nuclei Wash and Resuspension Buffer**

10m